

MODIFIED CEA NUCLEIC ACID AND EXPRESSION VECTORS

FIELD OF THE INVENTION

The present invention relates to a nucleic acid encoding a polypeptide and the use of the nucleic acid or polypeptide in preventing and / or treating cancer. In particular, the invention relates to improved vectors for the insertion and expression of foreign genes encoding tumor antigens for use in immunotherapeutic treatment of cancer.

BACKGROUND OF THE INVENTION

There has been tremendous increase in last few years in the development of cancer vaccines with Tumour-associated antigens (TAAs) due to the great advances in identification of molecules based on the expression profiling on primary tumours and normal cells with the help of several techniques such as high density microarray, SEREX, immunohistochemistry (IHC), RT-PCR, in-situ hybridization (ISH) and laser capture microscopy (Rosenberg, Immunity, 1999; Sgroi et al, 1999, Schena et al, 1995, Offringa et al, 2000). The TAAs are antigens expressed or over-expressed by tumour cells and could be specific to one or several tumours for example CEA antigen is expressed in colorectal, breast and lung cancers. Sgroi et al (1999) identified several genes differentially expressed in invasive and metastatic carcinoma cells with combined use of laser capture microdissection and cDNA microarrays. Several delivery systems like DNA or viruses could be used for therapeutic vaccination against human cancers (Bonnet et al, 2000) and can elicit immune responses and also break immune tolerance against TAAs. Tumour cells can be rendered more immunogenic by inserting transgenes encoding T cell co-stimulatory molecules such as B7.1 or cytokines IFNgamma, IL2, GM-CSF etc. Co-expression of a TAA and a cytokine or a co-stimulatory molecule can develop effective therapeutic vaccine (Hodge et al, 95, Bronte et al, 1995, Chamberlain et al, 1996).

There is a need in the art for reagents and methodologies useful in stimulating an immune response to prevent or treat cancers. The present inventions provides such reagents and methodologies which overcome many of the difficulties encountered by others in attempting to treat cancers such as cancer. In particular, the present invention provides a novel coding sequence for CEA. This nucleotide sequence, CEA(6D)-1,2, includes sequence modifications that eliminate the expression of truncated forms of CEA as expressed from expression vectors. Such a modified sequence is desired by those of skill in the art to improve expression and immunization protocols for CEA.

SUMMARY OF THE INVENTION

The present invention provides an immunogenic target for administration to a patient to prevent and / or treat cancer. In particular, the immunogenic target is a CEA tumor antigen ("TA") and / or an angiogenesis-associated antigen ("AA"). In one embodiment, the immunogenic target is encoded by a modified CEA nucleotide sequence (CEA(6D)-1,2) that improves CEA expression in transfected cells. In certain embodiments, the TA and / or AA are administered to a patient as a nucleic acid contained within a plasmid or other delivery vector, such as a recombinant virus. The TA and / or AA may also be administered in combination with an immune stimulator, such as a co-stimulatory molecule or adjuvant.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. A. Illustration of plasmid p3'H6MCEA comprising the CEA coding sequence with the 6D modification under the control of partial H6 promoter. B. Illustration of plasmid pSE1544.9 (pUC18-mCEA repeat 1).

Figure 2. Illustration of plasmid pSE1616.44 (pUC18-mCEA-modified repeat 1).

Figure 3. Illustration of plasmid pSE1658.15 (p3'H6MCEA-modified repeat 1).

Figure 4. Illustration of plasmid pBSmCEA.

Figure 5. Illustration of plasmid pSE1686.1 (pUC18 mCEA modified repeat 2).

Figure 6. Illustration of plasmid pSE1696.1 (pUC18 mCEA modified repeat 2).

Figure 7. Illustration of plasmid p3'H6modMCEA-1st&2nd repeats.

Figure 8. Illustration of plasmid pNVQH6MCEA(6D1st&2nd).

Figure 9A-D. Comparison of nucleotide sequence of CAP(6D) and CAP(6D)-1,2. Differences between the sequences are underlined.

Figure 10. PCR analysis to confirm the presence of CAP(6D)-1,2 in NYVAC DNA.

Figure 11. Immunoblot illustrating the lack of truncated CEA in cells expressing CAP(6D)-1,2.

DETAILED DESCRIPTION

The present invention provides reagents and methodologies useful for treating and / or preventing cancer. All references cited within this application are incorporated by reference.

In one embodiment, the present invention relates to the induction or enhancement of an immune response against one or more tumor antigens ("TA") to prevent and / or treat cancer. In certain embodiments, one or more TAs may be combined. In preferred embodiments, the immune response results from expression of a TA in a host cell following administration of a nucleic acid vector encoding the tumor antigen or the tumor antigen itself in the form of a peptide or polypeptide, for example.

As used herein, an "antigen" is a molecule (such as a polypeptide) or a portion thereof that produces an immune response in a host to whom the antigen has been administered. The immune response may include the production of antibodies that bind to at least one epitope of the antigen and / or the generation of a cellular immune response against cells expressing an epitope of the antigen. The response may be an enhancement of a current immune response by, for example, causing increased antibody production, production of antibodies with increased affinity for the antigen, or an increased cellular response (i.e., increased T cells). An antigen that produces an immune response may alternatively be referred to as being immunogenic or as an immunogen. In describing the present invention, a TA may be referred to as an "immunogenic target".

TA includes both tumor-associated antigens (TAAs) and tumor-specific antigens (TSAs), where a cancerous cell is the source of the antigen. A TAA is an antigen that is expressed on the surface of a tumor cell in higher amounts than is observed on normal cells or an antigen that is expressed on normal cells during fetal development. A TSA is an antigen that is unique to tumor cells and is not expressed on normal cells. TA further includes TAAs or TSAs, antigenic fragments thereof, and modified versions that retain their antigenicity.

TAs are typically classified into five categories according to their expression pattern, function, or genetic origin: cancer-testis (CT) antigens (i.e., MAGE, NY-ESO-1); melanocyte differentiation antigens (i.e., Melan A/MART-1, tyrosinase, gp100); mutational antigens (i.e., MUC-1, p53, CDK-4); overexpressed 'self' antigens (i.e., HER-2/neu, p53); and, viral antigens (i.e., HPV, EBV). For the purposes of practicing the present invention, a suitable TA is any TA that induces or enhances an anti-tumor immune response in a host to whom the TA has been administered. Suitable TAs include, for example, gp100 (Cox et al., *Science*, 264:716-719 (1994)), MART-1/Melan A (Kawakami et al., *J. Exp. Med.*, 180:347-352 (1994)), gp75 (TRP-1) (Wang et al., *J. Exp. Med.*, 186:1131-1140 (1996)), tyrosinase (Wolfel et al., *Eur. J. Immunol.*, 24:759-764 (1994); WO 200175117; WO 200175016; WO

200175007), NY-ESO-1 (WO 98/14464; WO 99/18206), melanoma proteoglycan (Hellstrom et al., *J. Immunol.*, 130:1467-1472 (1983)), MAGE family antigens (i.e., MAGE-1, 2,3,4,6,12, 51; Van der Bruggen et al., *Science*, 254:1643-1647 (1991); U.S. Pat. Nos. 6,235,525; CN 1319611), BAGE family antigens (Boel et al., *Immunity*, 2:167-175 (1995)),

5 GAGE family antigens (i.e., GAGE-1,2; Van den Eynde et al., *J. Exp. Med.*, 182:689-698 (1995); U.S. Pat. No. 6,013,765), RAGE family antigens (i.e., RAGE-1; Gaugler et al., *Immunogenetics*, 44:323-330 (1996); U.S. Pat. No. 5,939,526), N-acetylglucosaminyltransferase-V (Guilloux et al., *J. Exp. Med.*, 183:1173-1183 (1996)), p15 (Robbins et al., *J. Immunol.* 154:5944-5950 (1995)), β -catenin (Robbins et al., *J. Exp. Med.*,

10 183:1185-1192 (1996)), MUM-1 (Coulie et al., *Proc. Natl. Acad. Sci. USA*, 92:7976-7980 (1995)), cyclin dependent kinase-4 (CDK4) (Wolfel et al., *Science*, 269:1281-1284 (1995)), p21-ras (Fossum et al., *Int. J. Cancer*, 56:40-45 (1994)), BCR-*abl* (Bocchia et al., *Blood*, 85:2680-2684 (1995)), p53 (Theobald et al., *Proc. Natl. Acad. Sci. USA*, 92:11993-11997 (1995)), p185 HER2/neu (erb-B1; Fisk et al., *J. Exp. Med.*, 181:2109-2117 (1995)),

15 epidermal growth factor receptor (EGFR) (Harris et al., *Breast Cancer Res. Treat*, 29:1-2 (1994)), carcinoembryonic antigens (CEA) (Kwong et al., *J. Natl. Cancer Inst.*, 85:982-990 (1995) U.S. Pat. Nos. 5,756,103; 5,274,087; 5,571,710; 6,071,716; 5,698,530; 6,045,802; EP 263933; EP 346710; and, EP 784483); carcinoma-associated mutated mucins (i.e., MUC-1 gene products; Jerome et al., *J. Immunol.*, 151:1654-1662 (1993)); EBNA gene products of

20 EBV (i.e., EBNA-1; Rickinson et al., *Cancer Surveys*, 13:53-80 (1992)); E7, E6 proteins of human papillomavirus (Ressing et al., *J. Immunol.*, 154:5934-5943 (1995)); prostate specific antigen (PSA; Xue et al., *The Prostate*, 30:73-78 (1997)); prostate specific membrane antigen (PSMA; Israeli, et al., *Cancer Res.*, 54:1807-1811 (1994)); idiotypic epitopes or antigens, for example, immunoglobulin idiotypes or T cell receptor idiotypes (Chen et al., *J.*

25 *Immunol.*, 153:4775-4787 (1994)); KSA (U.S. Patent No. 5,348,887), kinesin 2 (Dietz, et al. *Biochem Biophys Res Commun* 2000 Sep 7;275(3):731-8), HIP-55, TGF β -1 anti-apoptotic factor (Toomey, et al. *Br J Biomed Sci* 2001;58(3):177-83), tumor protein D52 (Bryne J.A., et al., *Genomics*, 35:523-532 (1996)), H1FT, NY-BR-1 (WO 01/47959), NY-BR-62, NY-BR-75, NY-BR-85, NY-BR-87, NY-BR-96 (Scanlan, M. *Serologic and Bioinformatic*

30 *Approaches to the Identification of Human Tumor Antigens*, in *Cancer Vaccines 2000*, Cancer Research Institute, New York, NY), AAC2-1, or AAC2-2, including "wild-type" (i.e., normally encoded by the genome, naturally-occurring), modified, and mutated versions as

well as other fragments and derivatives thereof. Any of these TAs may be utilized alone or in combination with one another in a co-immunization protocol.

In certain cases, it may be beneficial to co-immunize patients with both TA and other antigens, such as angiogenesis-associated antigens ("AA"). An AA is an immunogenic molecule (i.e., peptide, polypeptide) associated with cells involved in the induction and / or continued development of blood vessels. For example, an AA may be expressed on an endothelial cell ("EC"), which is a primary structural component of blood vessels. Where the cancer is cancer, it is preferred that the AA be found within or near blood vessels that supply a tumor. Immunization of a patient against an AA preferably results in an anti-AA immune response whereby angiogenic processes that occur near or within tumors are prevented and / or inhibited.

Exemplary AAs include, for example, vascular endothelial growth factor (i.e., VEGF; Bernardini, et al. *J. Urol.*, 2001, 166(4): 1275-9; Starnes, et al. *J. Thorac. Cardiovasc. Surg.*, 2001, 122(3): 518-23), the VEGF receptor (i.e., VEGF-R, flk-1/KDR; Starnes, et al. *J. Thorac. Cardiovasc. Surg.*, 2001, 122(3): 518-23), EPH receptors (i.e., EPHA2; Gerety, et al. 1999, *Cell*, 4: 403-414), epidermal growth factor receptor (i.e., EGFR; Ciardeillo, et al. *Clin. Cancer Res.*, 2001, 7(10): 2958-70), basic fibroblast growth factor (i.e., bFGF; Davidson, et al. *Clin. Exp. Metastasis* 2000, 18(6): 501-7; Poon, et al. *Am J. Surg.*, 2001, 182(3):298-304), platelet-derived cell growth factor (i.e., PDGF-B), platelet-derived endothelial cell growth factor (PD-ECGF; Hong, et al. *J. Mol. Med.*, 2001, 8(2):141-8), transforming growth factors (i.e., TGF- α ; Hong, et al. *J. Mol. Med.*, 2001, 8(2):141-8), endoglin (Balza, et al. *Int. J. Cancer*, 2001, 94: 579-585), Id proteins (Benezra, R. *Trends Cardiovasc. Med.*, 2001, 11(6):237-41), proteases such as uPA, uPAR, and matrix metalloproteinases (MMP-2, MMP-9; Djonov, et al. *J. Pathol.*, 2001, 195(2):147-55), nitric oxide synthase (*Am. J. Ophthalmol.*, 2001, 132(4):551-6), aminopeptidase (Rouslhati, E. *Nature Cancer*, 2: 84-90, 2002), thrombospondins (i.e., TSP-1, TSP-2; Alvarez, et al. *Gynecol. Oncol.*, 2001, 82(2):273-8; Seki, et al. *Int. J. Oncol.*, 2001, 19(2):305-10), k-ras (Zhang, et al. *Cancer Res.*, 2001, 61(16):6050-4), Wnt (Zhang, et al. *Cancer Res.*, 2001, 61(16):6050-4), cyclin-dependent kinases (CDKs; *Drug Resist. Updat.* 2000, 3(2):83-88), microtubules (Timar, et al. 2001. *Path. Oncol. Res.*, 7(2): 85-94), heat shock proteins (i.e., HSP90 (Timar, *supra*)), heparin-binding factors (i.e., heparinase; Gohji, et al. *Int. J. Cancer*, 2001, 95(5):295-301), synthases (i.e., ATP synthase, thymidilate synthase), collagen receptors, integrins (i.e., $\alpha\upsilon\beta 3$, $\alpha\upsilon\beta 5$, $\alpha 1\beta 1$, $\alpha 2\beta 1$, $\alpha 5\beta 1$), the surface proteoglycan NG2, AAC2-1 (SEQ ID NO.:1), or AAC2-2

(SEQ ID NO.:2), among others, including “wild-type” (i.e., normally encoded by the genome, naturally-occurring), modified, mutated versions as well as other fragments and derivatives thereof. Any of these targets may be suitable in practicing the present invention, either alone or in combination with one another or with other agents.

5 In certain embodiments, a nucleic acid molecule encoding an immunogenic target is utilized. The nucleic acid molecule may comprise or consist of a nucleotide sequence encoding one or more immunogenic targets, or fragments or derivatives thereof, such as that contained in a DNA insert in an ATCC Deposit. The term “nucleic acid sequence” or “nucleic acid molecule” refers to a DNA or RNA sequence. The term encompasses
10 molecules formed from any of the known base analogs of DNA and RNA such as, but not limited to 4-acetylcytosine, 8-hydroxy-N6-methyladenosine, aziridinyl-cytosine, pseudoisocytosine, 5-(carboxyhydroxymethyl) uracil, 5-fluorouracil, 5-bromouracil, 5-carboxymethylaminomethyl-2-thiouracil, 5-carboxy-methylaminomethyluracil, dihydrouracil, inosine, N6-iso-pentenyladenine, 1-methyladenine, 1-methylpseudouracil, 1-
15 methylguanine, 1-methylinosine, 2,2-dimethyl-guanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-methyladenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyamino-methyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarbonyl-methyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, oxybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, N-
20 uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, pseudouracil, queosine, 2-thiocytosine, and 2,6-diaminopurine, among others.

An isolated nucleic acid molecule is one that: (1) is separated from at least about 50 percent of proteins, lipids, carbohydrates, or other materials with which it is naturally found
25 when total nucleic acid is isolated from the source cells; (2) is not be linked to all or a portion of a polynucleotide to which the nucleic acid molecule is linked in nature; (3) is operably linked to a polynucleotide which it is not linked to in nature; and / or, (4) does not occur in nature as part of a larger polynucleotide sequence. Preferably, the isolated nucleic acid molecule of the present invention is substantially free from any other contaminating nucleic
30 acid molecule(s) or other contaminants that are found in its natural environment that would interfere with its use in polypeptide production or its therapeutic, diagnostic, prophylactic or research use. As used herein, the term “naturally occurring” or “native” or “naturally found” when used in connection with biological materials such as nucleic acid molecules,

polypeptides, host cells, and the like, refers to materials which are found in nature and are not manipulated by man. Similarly, “non-naturally occurring” or “non-native” as used herein refers to a material that is not found in nature or that has been structurally modified or synthesized by man.

5 The identity of two or more nucleic acid or polypeptide molecules is determined by comparing the sequences. As known in the art, “identity” means the degree of sequence relatedness between nucleic acid molecules or polypeptides as determined by the match between the units making up the molecules (i.e., nucleotides or amino acid residues). Identity measures the percent of identical matches between the smaller of two or more sequences with
10 gap alignments (if any) addressed by a particular mathematical model or computer program (i.e., an algorithm). Identity between nucleic acid sequences may also be determined by the ability of the related sequence to hybridize to the nucleic acid sequence or isolated nucleic acid molecule. In defining such sequences, the term “highly stringent conditions” and “moderately stringent conditions” refer to procedures that permit hybridization of nucleic
15 acid strands whose sequences are complementary, and to exclude hybridization of significantly mismatched nucleic acids. Examples of “highly stringent conditions” for hybridization and washing are 0.015 M sodium chloride, 0.0015 M sodium citrate at 65-68°C or 0.015 M sodium chloride, 0.0015 M sodium citrate, and 50% formamide at 42°C. (see, for example, Sambrook, Fritsch & Maniatis, *Molecular Cloning: A Laboratory Manual* (2nd ed.,
20 Cold Spring Harbor Laboratory, 1989); Anderson *et al.*, *Nucleic Acid Hybridisation: A Practical Approach* Ch. 4 (IRL Press Limited)). The term “moderately stringent conditions” refers to conditions under which a DNA duplex with a greater degree of base pair mismatching than could occur under “highly stringent conditions” is able to form. Exemplary moderately stringent conditions are 0.015 M sodium chloride, 0.0015 M sodium
25 citrate at 50-65°C or 0.015 M sodium chloride, 0.0015 M sodium citrate, and 20% formamide at 37-50°C. By way of example, moderately stringent conditions of 50°C in 0.015 M sodium ion will allow about a 21% mismatch. During hybridization, other agents may be included in the hybridization and washing buffers for the purpose of reducing non-specific and/or background hybridization. Examples are 0.1% bovine serum albumin, 0.1% polyvinyl-
30 pyrrolidone, 0.1% sodium pyrophosphate, 0.1% sodium dodecylsulfate, NaDodSO₄, (SDS), ficoll, Denhardt’s solution, sonicated salmon sperm DNA (or another non-complementary DNA), and dextran sulfate, although other suitable agents can also be used. The concentration and types of these additives can be changed without substantially affecting the

stringency of the hybridization conditions. Hybridization experiments are usually carried out at pH 6.8-7.4; however, at typical ionic strength conditions, the rate of hybridization is nearly independent of pH.

In preferred embodiments of the present invention, vectors are used to transfer a nucleic acid sequence encoding a polypeptide to a cell. A vector is any molecule used to transfer a nucleic acid sequence to a host cell. In certain cases, an expression vector is utilized. An expression vector is a nucleic acid molecule that is suitable for transformation of a host cell and contains nucleic acid sequences that direct and / or control the expression of the transferred nucleic acid sequences. Expression includes, but is not limited to, processes such as transcription, translation, and splicing, if introns are present. Expression vectors typically comprise one or more flanking sequences operably linked to a heterologous nucleic acid sequence encoding a polypeptide. Flanking sequences may be homologous (i.e., from the same species and / or strain as the host cell), heterologous (i.e., from a species other than the host cell species or strain), hybrid (i.e., a combination of flanking sequences from more than one source), or synthetic, for example.

A flanking sequence is preferably capable of effecting the replication, transcription and / or translation of the coding sequence and is operably linked to a coding sequence. As used herein, the term operably linked refers to a linkage of polynucleotide elements in a functional relationship. For instance, a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the coding sequence. However, a flanking sequence need not necessarily be contiguous with the coding sequence, so long as it functions correctly. Thus, for example, intervening untranslated yet transcribed sequences can be present between a promoter sequence and the coding sequence and the promoter sequence may still be considered operably linked to the coding sequence. Similarly, an enhancer sequence may be located upstream or downstream from the coding sequence and affect transcription of the sequence.

In certain embodiments, it is preferred that the flanking sequence is a transcriptional regulatory region that drives high-level gene expression in the target cell. The transcriptional regulatory region may comprise, for example, a promoter, enhancer, silencer, repressor element, or combinations thereof. The transcriptional regulatory region may be either constitutive, tissue-specific, cell-type specific (i.e., the region drives higher levels of transcription in a one type of tissue or cell as compared to another), or regulatable (i.e., responsive to interaction with a compound such as tetracycline). The source of a

transcriptional regulatory region may be any prokaryotic or eukaryotic organism, any vertebrate or invertebrate organism, or any plant, provided that the flanking sequence functions in a cell by causing transcription of a nucleic acid within that cell. A wide variety of transcriptional regulatory regions may be utilized in practicing the present invention.

5 Suitable transcriptional regulatory regions include the CMV promoter (i.e., the CMV-immediate early promoter); promoters from eukaryotic genes (i.e., the estrogen-inducible chicken ovalbumin gene, the interferon genes, the gluco-corticoid-inducible tyrosine aminotransferase gene, and the thymidine kinase gene); and the major early and late adenovirus gene promoters; the SV40 early promoter region (Bernoist and Chambon, 1981,
10 *Nature* 290:304-10); the promoter contained in the 3' long terminal repeat (LTR) of Rous sarcoma virus (RSV) (Yamamoto, *et al.*, 1980, *Cell* 22:787-97); the herpes simplex virus thymidine kinase (HSV-TK) promoter (Wagner *et al.*, 1981, *Proc. Natl. Acad. Sci. U.S.A.* 78:1444-45); the regulatory sequences of the metallothionine gene (Brinster *et al.*, 1982, *Nature* 296:39-42); prokaryotic expression vectors such as the beta-lactamase promoter
15 (Villa-Kamaroff *et al.*, 1978, *Proc. Natl. Acad. Sci. U.S.A.*, 75:3727-31); or the tac promoter (DeBoer *et al.*, 1983, *Proc. Natl. Acad. Sci. U.S.A.*, 80:21-25). Tissue- and / or cell-type specific transcriptional control regions include, for example, the elastase I gene control region which is active in pancreatic acinar cells (Swift *et al.*, 1984, *Cell* 38:639-46; Ornitz *et al.*,
20 1986, *Cold Spring Harbor Symp. Quant. Biol.* 50:399-409 (1986); MacDonald, 1987, *Hepatology* 7:425-515); the insulin gene control region which is active in pancreatic beta cells (Hanahan, 1985, *Nature* 315:115-22); the immunoglobulin gene control region which is active in lymphoid cells (Grosschedl *et al.*, 1984, *Cell* 38:647-58; Adames *et al.*, 1985, *Nature* 318:533-38; Alexander *et al.*, 1987, *Mol. Cell. Biol.*, 7:1436-44); the mouse mammary tumor virus control region in testicular, breast, lymphoid and mast cells (Leder *et al.*,
25 1986, *Cell* 45:485-95); the albumin gene control region in liver (Pinkert *et al.*, 1987, *Genes and Devel.* 1:268-76); the alpha-feto-protein gene control region in liver (Krumlauf *et al.*, 1985, *Mol. Cell. Biol.*, 5:1639-48; Hammer *et al.*, 1987, *Science* 235:53-58); the alpha 1-antitrypsin gene control region in liver (Kelsey *et al.*, 1987, *Genes and Devel.* 1:161-71); the beta-globin gene control region in myeloid cells (Mogam *et al.*, 1985, *Nature* 315:338-40; Kollias *et al.*,
30 1986, *Cell* 46:89-94); the myelin basic protein gene control region in oligodendrocyte cells in the brain (Readhead *et al.*, 1987, *Cell* 48:703-12); the myosin light chain-2 gene control region in skeletal muscle (Sani, 1985, *Nature* 314:283-86); the gonadotropic releasing hormone gene control region in the hypothalamus (Mason *et al.*,

1986, *Science* 234:1372-78), and the tyrosinase promoter in melanoma cells (Hart, I. Semin Oncol 1996 Feb;23(1):154-8; Siders, et al. Cancer Gene Ther 1998 Sep-Oct;5(5):281-91), among others. Other suitable promoters are known in the art.

As described above, enhancers may also be suitable flanking sequences. Enhancers are cis-acting elements of DNA, usually about 10-300 bp in length, that act on the promoter to increase transcription. Enhancers are typically orientation- and position-independent, having been identified both 5' and 3' to controlled coding sequences. Several enhancer sequences available from mammalian genes are known (i.e., globin, elastase, albumin, alpha-feto-protein and insulin). Similarly, the SV40 enhancer, the cytomegalovirus early promoter enhancer, the polyoma enhancer, and adenovirus enhancers are useful with eukaryotic promoter sequences. While an enhancer may be spliced into the vector at a position 5' or 3' to nucleic acid coding sequence, it is typically located at a site 5' from the promoter. Other suitable enhancers are known in the art, and would be applicable to the present invention.

While preparing reagents of the present invention, cells may need to be transfected or transformed. Transfection refers to the uptake of foreign or exogenous DNA by a cell, and a cell has been transfected when the exogenous DNA has been introduced inside the cell membrane. A number of transfection techniques are well known in the art (i.e., Graham *et al.*, 1973, *Virology* 52:456; Sambrook *et al.*, *Molecular Cloning, A Laboratory Manual* (Cold Spring Harbor Laboratories, 1989); Davis *et al.*, *Basic Methods in Molecular Biology* (Elsevier, 1986); and Chu *et al.*, 1981, *Gene* 13:197). Such techniques can be used to introduce one or more exogenous DNA moieties into suitable host cells.

In certain embodiments, it is preferred that transfection of a cell results in transformation of that cell. A cell is transformed when there is a change in a characteristic of the cell, being transformed when it has been modified to contain a new nucleic acid. Following transfection, the transfected nucleic acid may recombine with that of the cell by physically integrating into a chromosome of the cell, may be maintained transiently as an episomal element without being replicated, or may replicate independently as a plasmid. A cell is stably transformed when the nucleic acid is replicated with the division of the cell.

The present invention further provides isolated immunogenic targets in polypeptide form. A polypeptide is considered isolated where it: (1) has been separated from at least about 50 percent of polynucleotides, lipids, carbohydrates, or other materials with which it is naturally found when isolated from the source cell; (2) is not linked (by covalent or noncovalent interaction) to all or a portion of a polypeptide to which the "isolated

polypeptide" is linked in nature; (3) is operably linked (by covalent or noncovalent interaction) to a polypeptide with which it is not linked in nature; or, (4) does not occur in nature. Preferably, the isolated polypeptide is substantially free from any other contaminating polypeptides or other contaminants that are found in its natural environment that would interfere with its therapeutic, diagnostic, prophylactic or research use.

Immunogenic target polypeptides may be mature polypeptides, as defined herein, and may or may not have an amino terminal methionine residue, depending on the method by which they are prepared. Further contemplated are related polypeptides such as, for example, fragments, variants (i.e., allelic, splice), orthologs, homologues, and derivatives, for example, that possess at least one characteristic or activity (i.e., activity, antigenicity) of the immunogenic target. Also related are peptides, which refers to a series of contiguous amino acid residues having a sequence corresponding to at least a portion of the polypeptide from which its sequence is derived. In preferred embodiments, the peptide comprises about 5-10 amino acids, 10-15 amino acids, 15-20 amino acids, 20-30 amino acids, or 30-50 amino acids. In a more preferred embodiment, a peptide comprises 9-12 amino acids, suitable for presentation upon Class I MHC molecules, for example.

A fragment of a nucleic acid or polypeptide comprises a truncation of the sequence (i.e., nucleic acid or polypeptide) at the amino terminus (with or without a leader sequence) and / or the carboxy terminus. Fragments may also include variants (i.e., allelic, splice), orthologs, homologues, and other variants having one or more amino acid additions or substitutions or internal deletions as compared to the parental sequence. In preferred embodiments, truncations and/or deletions comprise about 10 amino acids, 20 amino acids, 30 amino acids, 40 amino acids, 50 amino acids, or more. The polypeptide fragments so produced will comprise about 10 amino acids, 25 amino acids, 30 amino acids, 40 amino acids, 50 amino acids, 60 amino acids, 70 amino acids, or more. Such polypeptide fragments may optionally comprise an amino terminal methionine residue. It will be appreciated that such fragments can be used, for example, to generate antibodies or cellular immune responses to immunogenic target polypeptides.

A variant is a sequence having one or more sequence substitutions, deletions, and/or additions as compared to the subject sequence. Variants may be naturally occurring or artificially constructed. Such variants may be prepared from the corresponding nucleic acid molecules. In preferred embodiments, the variants have from 1 to 3, or from 1 to 5, or from 1

to 10, or from 1 to 15, or from 1 to 20, or from 1 to 25, or from 1 to 30, or from 1 to 40, or from 1 to 50, or more than 50 amino acid substitutions, insertions, additions and/or deletions.

An allelic variant is one of several possible naturally-occurring alternate forms of a gene occupying a given locus on a chromosome of an organism or a population of organisms.

5 A splice variant is a polypeptide generated from one of several RNA transcript resulting from splicing of a primary transcript. An ortholog is a similar nucleic acid or polypeptide sequence from another species. For example, the mouse and human versions of an immunogenic target polypeptide may be considered orthologs of each other. A derivative of a sequence is one that is derived from a parental sequence those sequences having
10 substitutions, additions, deletions, or chemically modified variants. Variants may also include fusion proteins, which refers to the fusion of one or more first sequences (such as a peptide) at the amino or carboxy terminus of at least one other sequence (such as a heterologous peptide).

“Similarity” is a concept related to identity, except that similarity refers to a measure
15 of relatedness which includes both identical matches and conservative substitution matches. If two polypeptide sequences have, for example, 10/20 identical amino acids, and the remainder are all non-conservative substitutions, then the percent identity and similarity would both be 50%. If in the same example, there are five more positions where there are conservative substitutions, then the percent identity remains 50%, but the percent similarity
20 would be 75% (15/20). Therefore, in cases where there are conservative substitutions, the percent similarity between two polypeptides will be higher than the percent identity between those two polypeptides.

Substitutions may be conservative, or non-conservative, or any combination thereof. Conservative amino acid modifications to the sequence of a polypeptide (and the
25 corresponding modifications to the encoding nucleotides) may produce polypeptides having functional and chemical characteristics similar to those of a parental polypeptide. For example, a “conservative amino acid substitution” may involve a substitution of a native amino acid residue with a non-native residue such that there is little or no effect on the size, polarity, charge, hydrophobicity, or hydrophilicity of the amino acid residue at that position
30 and, in particular, does not result in decreased immunogenicity. Suitable conservative amino acid substitutions are shown in **Table I**.

Table I

Original Residues	Exemplary Substitutions	Preferred Substitutions
Ala	Val, Leu, Ile	Val
Arg	Lys, Gln, Asn	Lys
Asn	Gln	Gln
Asp	Glu	Glu
Cys	Ser, Ala	Ser
Gln	Asn	Asn
Glu	Asp	Asp
Gly	Pro, Ala	Ala
His	Asn, Gln, Lys, Arg	Arg
Ile	Leu, Val, Met, Ala, Phe, Norleucine	Leu
Leu	Norleucine, Ile, Val, Met, Ala, Phe	Ile
Lys	Arg, 1,4 Diamino-butyric Acid, Gln, Asn	Arg
Met	Leu, Phe, Ile	Leu
Phe	Leu, Val, Ile, Ala, Tyr	Leu
Pro	Ala	Gly
Ser	Thr, Ala, Cys	Thr
Thr	Ser	Ser
Trp	Tyr, Phe	Tyr
Tyr	Trp, Phe, Thr, Ser	Phe
Val	Ile, Met, Leu, Phe, Ala, Norleucine	Leu

A skilled artisan will be able to determine suitable variants of polypeptide using well-known techniques. For identifying suitable areas of the molecule that may be changed without destroying biological activity (i.e., MHC binding, immunogenicity), one skilled in the art may target areas not believed to be important for that activity. For example, when similar polypeptides with similar activities from the same species or from other species are known, one skilled in the art may compare the amino acid sequence of a polypeptide to such similar polypeptides. By performing such analyses, one can identify residues and portions of the molecules that are conserved among similar polypeptides. It will be appreciated that changes in areas of the molecule that are not conserved relative to such similar polypeptides would be less likely to adversely affect the biological activity and/or structure of a polypeptide. Similarly, the residues required for binding to MHC are known, and may be modified to improve binding. However, modifications resulting in decreased binding to MHC will not be appropriate in most situations. One skilled in the art would also know that, even in relatively conserved regions, one may substitute chemically similar amino acids for the naturally occurring residues while retaining activity. Therefore, even areas that may be important for biological activity or for structure may be subject to conservative amino acid

substitutions without destroying the biological activity or without adversely affecting the polypeptide structure.

Other preferred polypeptide variants include glycosylation variants wherein the number and/or type of glycosylation sites have been altered compared to the subject amino acid sequence. In one embodiment, polypeptide variants comprise a greater or a lesser number of N-linked glycosylation sites than the subject amino acid sequence. An N-linked glycosylation site is characterized by the sequence Asn-X-Ser or Asn-X-Thr, wherein the amino acid residue designated as X may be any amino acid residue except proline. The substitution of amino acid residues to create this sequence provides a potential new site for the addition of an N-linked carbohydrate chain. Alternatively, substitutions that eliminate this sequence will remove an existing N-linked carbohydrate chain. Also provided is a rearrangement of N-linked carbohydrate chains wherein one or more N-linked glycosylation sites (typically those that are naturally occurring) are eliminated and one or more new N-linked sites are created. To affect O-linked glycosylation of a polypeptide, one would modify serine and / or threonine residues.

Additional preferred variants include cysteine variants, wherein one or more cysteine residues are deleted or substituted with another amino acid (*e.g.*, serine) as compared to the subject amino acid sequence set. Cysteine variants are useful when polypeptides must be refolded into a biologically active conformation such as after the isolation of insoluble inclusion bodies. Cysteine variants generally have fewer cysteine residues than the native protein, and typically have an even number to minimize interactions resulting from unpaired cysteines.

In other embodiments, the isolated polypeptides of the current invention include fusion polypeptide segments that assist in purification of the polypeptides. Fusions can be made either at the amino terminus or at the carboxy terminus of the subject polypeptide variant thereof. Fusions may be direct with no linker or adapter molecule or may be through a linker or adapter molecule. A linker or adapter molecule may be one or more amino acid residues, typically from about 20 to about 50 amino acid residues. A linker or adapter molecule may also be designed with a cleavage site for a DNA restriction endonuclease or for a protease to allow for the separation of the fused moieties. It will be appreciated that once constructed, the fusion polypeptides can be derivatized according to the methods described herein. Suitable fusion segments include, among others, metal binding domains (*e.g.*, a poly-histidine segment), immunoglobulin binding domains (*i.e.*, Protein A, Protein G, T cell,

B cell, Fc receptor, or complement protein antibody-binding domains), sugar binding domains (e.g., a maltose binding domain), and/or a "tag" domain (i.e., at least a portion of α -galactosidase, a strep tag peptide, a T7 tag peptide, a FLAG peptide, or other domains that can be purified using compounds that bind to the domain, such as monoclonal antibodies).

5 This tag is typically fused to the polypeptide upon expression of the polypeptide, and can serve as a means for affinity purification of the sequence of interest polypeptide from the host cell. Affinity purification can be accomplished, for example, by column chromatography using antibodies against the tag as an affinity matrix. Optionally, the tag can subsequently be removed from the purified sequence of interest polypeptide by various means such as using
10 certain peptidases for cleavage. As described below, fusions may also be made between a TA and a co-stimulatory components such as the chemokines CXCL10 (IP-10), CCL7 (MCP-3), or CCL5 (RANTES), for example.

A fusion motif may enhance transport of an immunogenic target to an MHC processing compartment, such as the endoplasmic reticulum. These sequences, referred to as
15 transduction or transcytosis sequences, include sequences derived from HIV tat (see Kim et al. 1997 J. Immunol. 159:1666), *Drosophila* antennapedia (see Schutze-Redelmeier et al. 1996 J. Immunol. 157:650), or human period-1 protein (hPER1; in particular, SRRHHCRSKAKRSRHH).

In addition, the polypeptide or variant thereof may be fused to a homologous
20 polypeptide to form a homodimer or to a heterologous polypeptide to form a heterodimer. Heterologous peptides and polypeptides include, but are not limited to: an epitope to allow for the detection and/or isolation of a fusion polypeptide; a transmembrane receptor protein or a portion thereof, such as an extracellular domain or a transmembrane and intracellular domain; a ligand or a portion thereof which binds to a transmembrane receptor protein; an
25 enzyme or portion thereof which is catalytically active; a polypeptide or peptide which promotes oligomerization, such as a leucine zipper domain; a polypeptide or peptide which increases stability, such as an immunoglobulin constant region; and a polypeptide which has a therapeutic activity different from the polypeptide or variant thereof.

In certain embodiments, it may be advantageous to combine a nucleic acid sequence
30 encoding an immunogenic target, polypeptide, or derivative thereof with one or more co-stimulatory component(s) such as cell surface proteins, cytokines or chemokines in a composition of the present invention. The co-stimulatory component may be included in the composition as a polypeptide or as a nucleic acid encoding the polypeptide, for example.

Suitable co-stimulatory molecules include, for instance, polypeptides that bind members of the CD28 family (i.e., CD28, ICOS; Hutloff, et al. *Nature* 1999, 397: 263–265; Peach, et al. *J Exp Med* 1994, 180: 2049–2058) such as the CD28 binding polypeptides B7.1 (CD80; Schwartz, 1992; Chen et al, 1992; Ellis, et al. *J. Immunol.*, 156(8): 2700-9) and B7.2 (CD86; Ellis, et al. *J. Immunol.*, 156(8): 2700-9); polypeptides which bind members of the integrin family (i.e., LFA-1 (CD11a / CD18); Sedwick, et al. *J Immunol* 1999, 162: 1367–1375; Wülfing, et al. *Science* 1998, 282: 2266–2269; Lub, et al. *Immunol Today* 1995, 16: 479–483) including members of the ICAM family (i.e., ICAM-1, -2 or -3); polypeptides which bind CD2 family members (i.e., CD2, signalling lymphocyte activation molecule (CDw150 or “SLAM”; Aversa, et al. *J Immunol* 1997, 158: 4036–4044)) such as CD58 (LFA-3; CD2 ligand; Davis, et al. *Immunol Today* 1996, 17: 177–187) or SLAM ligands (Sayos, et al. *Nature* 1998, 395: 462–469); polypeptides which bind heat stable antigen (HSA or CD24; Zhou, et al. *Eur J Immunol* 1997, 27: 2524–2528); polypeptides which bind to members of the TNF receptor (TNFR) family (i.e., 4-1BB (CD137; Vinay, et al. *Semin Immunol* 1998, 10: 481–489), OX40 (CD134; Weinberg, et al. *Semin Immunol* 1998, 10: 471–480; Higgins, et al. *J Immunol* 1999, 162: 486–493), and CD27 (Lens, et al. *Semin Immunol* 1998, 10: 491–499)) such as 4-1BBL (4-1BB ligand; Vinay, et al. *Semin Immunol* 1998, 10: 481–48; DeBenedette, et al. *J Immunol* 1997, 158: 551–559), TNFR associated factor-1 (TRAF-1; 4-1BB ligand; Saoulli, et al. *J Exp Med* 1998, 187: 1849–1862, Arch, et al. *Mol Cell Biol* 1998, 18: 558–565), TRAF-2 (4-1BB and OX40 ligand; Saoulli, et al. *J Exp Med* 1998, 187: 1849–1862; Oshima, et al. *Int Immunol* 1998, 10: 517–526, Kawamata, et al. *J Biol Chem* 1998, 273: 5808–5814), TRAF-3 (4-1BB and OX40 ligand; Arch, et al. *Mol Cell Biol* 1998, 18: 558–565; Jang, et al. *Biochem Biophys Res Commun* 1998, 242: 613–620; Kawamata S, et al. *J Biol Chem* 1998, 273: 5808–5814), OX40L (OX40 ligand; Gramaglia, et al. *J Immunol* 1998, 161: 6510–6517), TRAF-5 (OX40 ligand; Arch, et al. *Mol Cell Biol* 1998, 18: 558–565; Kawamata, et al. *J Biol Chem* 1998, 273: 5808–5814), and CD70 (CD27 ligand; Couderc, et al. *Cancer Gene Ther.*, 5(3): 163-75). CD154 (CD40 ligand or “CD40L”; Gurunathan, et al. *J. Immunol.*, 1998, 161: 4563-4571; Sine, et al. *Hum. Gene Ther.*, 2001, 12: 1091-1102) may also be suitable.

One or more cytokines may also be suitable co-stimulatory components or “adjuvants”, either as polypeptides or being encoded by nucleic acids contained within the compositions of the present invention (Parmiani, et al. *Immunol Lett* 2000 Sep 15; 74(1): 41-

4; Berzofsky, et al. *Nature Immunol.* 1: 209-219). Suitable cytokines include, for example, interleukin-2 (IL-2) (Rosenberg, et al. *Nature Med.* 4: 321-327 (1998)), IL-4, IL-7, IL-12 (reviewed by Pardoll, 1992; Harries, et al. *J. Gene Med.* 2000 Jul-Aug;2(4):243-9; Rao, et al. *J. Immunol.* 156: 3357-3365 (1996)), IL-15 (Xin, et al. *Vaccine*, 17:858-866, 1999), IL-16
5 (Cruikshank, et al. *J. Leuk Biol.* 67(6): 757-66, 2000), IL-18 (*J. Cancer Res. Clin. Oncol.* 2001. 127(12): 718-726), GM-CSF (CSF (Disis, et al. *Blood*, 88: 202-210 (1996)), tumor necrosis factor-alpha (TNF- α), or interferon-gamma (INF- γ). Other cytokines may also be suitable for practicing the present invention, as is known in the art.

Chemokines may also be utilized. For example, fusion proteins comprising CXCL10
10 (IP-10) and CCL7 (MCP-3) fused to a tumor self-antigen have been shown to induce anti-tumor immunity (Biragyn, et al. *Nature Biotech.* 1999, 17: 253-258). The chemokines CCL3 (MIP-1 α) and CCL5 (RANTES) (Boyer, et al. *Vaccine*, 1999, 17 (Supp. 2): S53-S64) may also be of use in practicing the present invention. Other suitable chemokines are known in the art.

15 It is also known in the art that suppressive or negative regulatory immune mechanisms may be blocked, resulting in enhanced immune responses. For instance, treatment with anti-CTLA-4 (Shrikant, et al. *Immunity*, 1996, 14: 145-155; Suttmuller, et al. *J. Exp. Med.*, 2001, 194: 823-832), anti-CD25 (Suttmuller, *supra*), anti-CD4 (Matsui, et al. *J. Immunol.*, 1999, 163: 184-193), the fusion protein IL13Ra2-Fc (Terabe, et al. *Nature Immunol.*, 2000, 1: 515-520), and combinations thereof (i.e., anti-CTLA-4 and anti-CD25,
20 Suttmuller, *supra*) have been shown to upregulate anti-tumor immune responses and would be suitable in practicing the present invention.

Any of these components may be used alone or in combination with other agents. For instance, it has been shown that a combination of CD80, ICAM-1 and LFA-3 ("TRICOM")
25 may potentiate anti-cancer immune responses (Hodge, et al. *Cancer Res.* 59: 5800-5807 (1999). Other effective combinations include, for example, IL-12 + GM-CSF (Ahlers, et al. *J. Immunol.*, 158: 3947-3958 (1997); Iwasaki, et al. *J. Immunol.* 158: 4591-4601 (1997)), IL-12 + GM-CSF + TNF- α (Ahlers, et al. *Int. Immunol.* 13: 897-908 (2001)), CD80 + IL-12 (Fruend, et al. *Int. J. Cancer*, 85: 508-517 (2000); Rao, et al. *supra*), and CD86 + GM-CSF +
30 IL-12 (Iwasaki, *supra*). One of skill in the art would be aware of additional combinations useful in carrying out the present invention. In addition, the skilled artisan would be aware of additional reagents or methods that may be used to modulate such mechanisms. These

reagents and methods, as well as others known by those of skill in the art, may be utilized in practicing the present invention.

Additional strategies for improving the efficiency of nucleic acid-based immunization may also be used including, for example, the use of self-replicating viral replicons (Caley, et al. 1999. *Vaccine*, 17: 3124-2135; Dubensky, et al. 2000. *Mol. Med.* 6: 723-732; Leitner, et al. 2000. *Cancer Res.* 60: 51-55), codon optimization (Liu, et al. 2000. *Mol. Ther.*, 1: 497-500; Dubensky, *supra*; Huang, et al. 2001. *J. Virol.* 75: 4947-4951), *in vivo* electroporation (Widera, et al. 2000. *J. Immunol.* 164: 4635-3640), incorporation of CpG stimulatory motifs (Gurunathan, et al. *Ann. Rev. Immunol.*, 2000, 18: 927-974; Leitner, *supra*), sequences for targeting of the endocytic or ubiquitin-processing pathways (Thomson, et al. 1998. *J. Virol.* 72: 2246-2252; Velders, et al. 2001. *J. Immunol.* 166: 5366-5373), prime-boost regimens (Gurunathan, *supra*; Sullivan, et al. 2000. *Nature*, 408: 605-609; Hanke, et al. 1998. *Vaccine*, 16: 439-445; Amara, et al. 2001. *Science*, 292: 69-74), and the use of mucosal delivery vectors such as *Salmonella* (Darji, et al. 1997. *Cell*, 91: 765-775; Woo, et al. 2001. *Vaccine*, 19: 2945-2954). Other methods are known in the art, some of which are described below.

Chemotherapeutic agents, radiation, anti-angiogenic compounds, or other agents may also be utilized in treating and / or preventing cancer using immunogenic targets (Sebti, et al. *Oncogene* 2000 Dec 27;19(56):6566-73). For example, in treating metastatic breast cancer, useful chemotherapeutic agents include cyclophosphamide, doxorubicin, paclitaxel, docetaxel, navelbine, capecitabine, and mitomycin C, among others. Combination chemotherapeutic regimens have also proven effective including cyclophosphamide + methotrexate + 5-fluorouracil; cyclophosphamide + doxorubicin + 5-fluorouracil; or, cyclophosphamide + doxorubicin, for example. Other compounds such as prednisone, a taxane, navelbine, mitomycin C, or vinblastine have been utilized for various reasons. A majority of breast cancer patients have estrogen-receptor positive (ER+) tumors and in these patients, endocrine therapy (i.e., tamoxifen) is preferred over chemotherapy. For such patients, tamoxifen or, as a second line therapy, progestins (medroxyprogesterone acetate or megestrol acetate) are preferred. Aromatase inhibitors (i.e., aminoglutethimide and analogs thereof such as letrozole) decrease the availability of estrogen needed to maintain tumor growth and may be used as second or third line endocrine therapy in certain patients.

Other cancers may require different chemotherapeutic regimens. For example, metastatic colorectal cancer is typically treated with Camptosar (irinotecan or CPT-11), 5-

fluorouracil or leucovorin, alone or in combination with one another. Proteinase and integrin inhibitors such as the MMP inhibitors marimastate (British Biotech), COL-3 (Collagenex), Neovastat (Aeterna), AG3340 (Agouron), BMS-275291 (Bristol Myers Squibb), CGS 27023A (Novartis) or the integrin inhibitors Vitaxin (Medimmune), or MED1522 (Merck KgaA) may also be suitable for use. As such, immunological targeting of immunogenic targets associated with colorectal cancer could be performed in combination with a treatment using those chemotherapeutic agents. Similarly, chemotherapeutic agents used to treat other types of cancers are well-known in the art and may be combined with the immunogenic targets described herein.

Many anti-angiogenic agents are known in the art and would be suitable for co-administration with the immunogenic target vaccines (see, for example, Timar, et al. 2001. *Pathology Oncol. Res.*, 7(2): 85-94). Such agents include, for example, physiological agents such as growth factors (i.e., ANG-2, NK1,2,4 (HGF), transforming growth factor beta (TGF- β)), cytokines (i.e., interferons such as IFN- α , - β , - γ , platelet factor 4 (PF-4), PR-39), proteases (i.e., cleaved AT-III, collagen XVIII fragment (Endostatin)), HmwKallikrein-d5 plasmin fragment (Angiostatin), prothrombin-FI-2, TSP-1), protease inhibitors (i.e., tissue inhibitor of metalloproteases such as TIMP-1, -2, or -3; maspin; plasminogen activator-inhibitors such as PAI-1; pigment epithelium derived factor (PEDF)), Tumstatin (available through ILEX, Inc.), antibody products (i.e., the collagen-binding antibodies HUIV26, HUI77, XL313; anti-VEGF; anti-integrin (i.e., Vitaxin, (Lxsys))), and glycosidases (i.e., heparinase-I, -III). "Chemical" or modified physiological agents known or believed to have anti-angiogenic potential include, for example, vinblastine, taxol, ketoconazole, thalidomide, dolestatin, combrestatin A, rapamycin (Guba, et al. 2002, *Nature Med.*, 8: 128-135), CEP-7055 (available from Cephalon, Inc.), flavone acetic acid, Bay 12-9566 (Bayer Corp.), AG3340 (Agouron, Inc.), CGS 27023A (Novartis), tetracycline derivatives (i.e., COL-3 (Collagenix, Inc.)), Neovastat (Aeterna), BMS-275291 (Bristol-Myers Squibb), low dose 5-FU, low dose methotrexate (MTX), irsofladine, radicicol, cyclosporine, captopril, celecoxib, D45152-sulphated polysaccharide, cationic protein (Protamine), cationic peptide-VEGF, Suramin (polysulphonated naphthyl urea), compounds that interfere with the function or production of VEGF (i.e., SU5416 or SU6668 (Sugen), PTK787/ZK22584 (Novartis)), Distamycin A, Angiozyme (ribozyme), isoflavinoids, staurosporine derivatives, genistein, EMD121974 (Merck KgaA), tyrphostins, isoquinolones, retinoic acid, carboxyamidotriazole, TNP-470, octreotide, 2-methoxyestradiol, aminosterols (i.e.,

squalamine), glutathione analogues (i.e., N-acetyl-L-cysteine), combretastatin A-4 (Oxigene), Eph receptor blocking agents (*Nature*, 414:933-938, 2001), Rh-Angiostatin, Rh-Endostatin (WO 01/93897), cyclic-RGD peptide, accutin-disintegrin, benzodiazepenes, humanized anti-avb3 Ab, Rh-PAI-2, amiloride, p-amidobenzamidine, anti-uPA ab, anti-uPAR Ab, L-phenylalanine-N-methylamides (i.e., Batimistat, Marimastat), AG3340, and minocycline. Many other suitable agents are known in the art and would suffice in practicing the present invention.

The present invention may also be utilized in combination with “non-traditional” methods of treating cancer. For example, it has recently been demonstrated that administration of certain anaerobic bacteria may assist in slowing tumor growth. In one study, *Clostridium novyi* was modified to eliminate a toxin gene carried on a phage episome and administered to mice with colorectal tumors (Dang, et al. *P.N.A.S. USA*, 98(26): 15155-15160, 2001). In combination with chemotherapy, the treatment was shown to cause tumor necrosis in the animals. The reagents and methodologies described in this application may be combined with such treatment methodologies.

Nucleic acids encoding immunogenic targets may be administered to patients by any of several available techniques. Various viral vectors that have been successfully utilized for introducing a nucleic acid to a host include retrovirus, adenovirus, adeno-associated virus (AAV), herpes virus, and poxvirus, among others. It is understood in the art that many such viral vectors are available in the art. The vectors of the present invention may be constructed using standard recombinant techniques widely available to one skilled in the art. Such techniques may be found in common molecular biology references such as *Molecular Cloning: A Laboratory Manual* (Sambrook, et al., 1989, Cold Spring Harbor Laboratory Press), *Gene Expression Technology* (Methods in Enzymology, Vol. 185, edited by D. Goeddel, 1991. Academic Press, San Diego, CA), and *PCR Protocols: A Guide to Methods and Applications* (Innis, et al. 1990. Academic Press, San Diego, CA).

Preferred retroviral vectors are derivatives of lentivirus as well as derivatives of murine or avian retroviruses. Examples of suitable retroviral vectors include, for example, Moloney murine leukemia virus (MoMuLV), Harvey murine sarcoma virus (HaMuSV), murine mammary tumor virus (MuMTV), SIV, BIV, HIV and Rous Sarcoma Virus (RSV). A number of retroviral vectors can incorporate multiple exogenous nucleic acid sequences. As recombinant retroviruses are defective, they require assistance in order to produce infectious vector particles. This assistance can be provided by, for example, helper cell lines

encoding retrovirus structural genes. Suitable helper cell lines include Ψ 2, PA317 and PA12, among others. The vector virions produced using such cell lines may then be used to infect a tissue cell line, such as NIH 3T3 cells, to produce large quantities of chimeric retroviral virions. Retroviral vectors may be administered by traditional methods (i.e., injection) or by
5 implantation of a "producer cell line" in proximity to the target cell population (Culver, K., *et al.*, 1994, *Hum. Gene Ther.*, 5 (3): 343-79; Culver, K., *et al.*, *Cold Spring Harb. Symp. Quant. Biol.*, 59: 685-90); Oldfield, E., 1993, *Hum. Gene Ther.*, 4 (1): 39-69). The producer cell line is engineered to produce a viral vector and releases viral particles in the vicinity of the target cell. A portion of the released viral particles contact the target cells and infect those
10 cells, thus delivering a nucleic acid of the present invention to the target cell. Following infection of the target cell, expression of the nucleic acid of the vector occurs.

Adenoviral vectors have proven especially useful for gene transfer into eukaryotic cells (Rosenfeld, M., *et al.*, 1991, *Science*, 252 (5004): 431-4; Crystal, R., *et al.*, 1994, *Nat. Genet.*, 8 (1): 42-51), the study eukaryotic gene expression (Levrero, M., *et al.*, 1991, *Gene*,
15 101 (2): 195-202), vaccine development (Graham, F. and Prevec, L., 1992, *Biotechnology*, 20: 363-90), and in animal models (Stratford-Perricaudet, L., *et al.*, 1992, *Bone Marrow Transplant.*, 9 (Suppl. 1): 151-2 ; Rich, D., *et al.*, 1993, *Hum. Gene Ther.*, 4 (4): 461-76). Experimental routes for administering recombinant Ad to different tissues *in vivo* have included intratracheal instillation (Rosenfeld, M., *et al.*, 1992, *Cell*, 68 (1): 143-55) injection
20 into muscle (Quantin, B., *et al.*, 1992, *Proc. Natl. Acad. Sci. U.S.A.*, 89 (7): 2581-4), peripheral intravenous injection (Herz, J., and Gerard, R., 1993, *Proc. Natl. Acad. Sci. U.S.A.*, 90 (7): 2812-6) and stereotactic inoculation to brain (Le Gal La Salle, G., *et al.*, 1993, *Science*, 259 (5097): 988-90), among others.

Adeno-associated virus (AAV) demonstrates high-level infectivity, broad host range
25 and specificity in integrating into the host cell genome (Hermonat, P., *et al.*, 1984, *Proc. Natl. Acad. Sci. U.S.A.*, 81 (20): 6466-70). And Herpes Simplex Virus type-1 (HSV-1) is yet another attractive vector system, especially for use in the nervous system because of its neurotropic property (Geller, A., *et al.*, 1991, *Trends Neurosci.*, 14 (10): 428-32; Glorioso, *et al.*, 1995, *Mol. Biotechnol.*, 4 (1): 87-99; Glorioso, *et al.*, 1995, *Annu. Rev. Microbiol.*, 49:
30 675-710).

Poxvirus is another useful expression vector (Smith, *et al.* 1983, *Gene*, 25 (1): 21-8; Moss, *et al.*, 1992, *Biotechnology*, 20: 345-62; Moss, *et al.*, 1992, *Curr. Top. Microbiol. Immunol.*, 158: 25-38; Moss, *et al.* 1991. *Science*, 252: 1662-1667). Poxviruses shown to be

useful include vaccinia, NYVAC, avipox, fowlpox, canarypox, ALVAC, and ALVAC(2), among others.

NYVAC (vP866) was derived from the Copenhagen vaccine strain of vaccinia virus by deleting six nonessential regions of the genome encoding known or potential virulence factors (see, for example, U.S. Pat. Nos. 5,364,773 and 5,494,807). The deletion loci were also engineered as recipient loci for the insertion of foreign genes. The deleted regions are: thymidine kinase gene (TK; J2R); hemorrhagic region (u; B13R+B14R); A type inclusion body region (ATI; A26L); hemagglutinin gene (HA; A56R); host range gene region (C7L-K1L); and, large subunit, ribonucleotide reductase (I4L). NYVAC is a genetically engineered vaccinia virus strain that was generated by the specific deletion of eighteen open reading frames encoding gene products associated with virulence and host range. NYVAC has been shown to be useful for expressing TAs (see, for example, U.S. Pat. No. 6,265,189). NYVAC (vP866), vP994, vCP205, vCP1433, placZH6H4Lreverse, pMPC6H6K3E3 and pC3H6FHVB were also deposited with the ATCC under the terms of the Budapest Treaty, accession numbers VR-2559, VR-2558, VR-2557, VR-2556, ATCC-97913, ATCC-97912, and ATCC-97914, respectively.

ALVAC-based recombinant viruses (i.e., ALVAC-1 and ALVAC-2) are also suitable for use in practicing the present invention (see, for example, U.S. Pat. No. 5,756,103). ALVAC(2) is identical to ALVAC(1) except that ALVAC(2) genome comprises the vaccinia E3L and K3L genes under the control of vaccinia promoters (U.S. Pat. No. 6,130,066; Beattie et al., 1995a, 1995b, 1991; Chang et al., 1992; Davies et al., 1993). Both ALVAC(1) and ALVAC(2) have been demonstrated to be useful in expressing foreign DNA sequences, such as TAs (Tartaglia et al., 1993 a,b; U.S. Pat. No. 5,833,975). ALVAC was deposited under the terms of the Budapest Treaty with the American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, Va. 20110-2209, USA, ATCC accession number VR-2547.

Another useful poxvirus vector is TROVAC. TROVAC refers to an attenuated fowlpox that was a plaque-cloned isolate derived from the FP-1 vaccine strain of fowlpoxvirus which is licensed for vaccination of 1 day old chicks. TROVAC was likewise deposited under the terms of the Budapest Treaty with the ATCC, accession number 2553.

“Non-viral” plasmid vectors may also be suitable in practicing the present invention. Preferred plasmid vectors are compatible with bacterial, insect, and / or mammalian host cells. Such vectors include, for example, PCR-II, pCR3, and pcDNA3.1 (Invitrogen, San Diego, CA), pBSII (Stratagene, La Jolla, CA), pET15 (Novagen, Madison, WI), pGEX

(Pharmacia Biotech, Piscataway, NJ), pEGFP-N2 (Clontech, Palo Alto, CA), pETL (BlueBacII, Invitrogen), pDSR-alpha (PCT pub. No. WO 90/14363) and pFastBacDual (Gibco-BRL, Grand Island, NY) as well as Bluescript[®] plasmid derivatives (a high copy number COLE1-based phagemid, Stratagene Cloning Systems, La Jolla, CA), PCR cloning plasmids designed for cloning Taq-amplified PCR products (e.g., TOPO[™] TA cloning[®] kit, PCR2.1[®] plasmid derivatives, Invitrogen, Carlsbad, CA). Bacterial vectors may also be used with the current invention. These vectors include, for example, *Shigella*, *Salmonella*, *Vibrio cholerae*, *Lactobacillus*, *Bacille calmette guérin (BCG)*, and *Streptococcus* (see for example, WO 88/6626; WO 90/0594; WO 91/13157; WO 92/1796; and WO 92/21376). Many other non-viral plasmid expression vectors and systems are known in the art and could be used with the current invention.

Suitable nucleic acid delivery techniques include DNA-ligand complexes, adenovirus-ligand-DNA complexes, direct injection of DNA, CaPO₄ precipitation, gene gun techniques, electroporation, and colloidal dispersion systems, among others. Colloidal dispersion systems include macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. The preferred colloidal system of this invention is a liposome, which are artificial membrane vesicles useful as delivery vehicles *in vitro* and *in vivo*. RNA, DNA and intact virions can be encapsulated within the aqueous interior and be delivered to cells in a biologically active form (Fraley, R., *et al.*, 1981, *Trends Biochem. Sci.*, 6: 77). The composition of the liposome is usually a combination of phospholipids, particularly high-phase-transition-temperature phospholipids, usually in combination with steroids, especially cholesterol. Other phospholipids or other lipids may also be used. The physical characteristics of liposomes depend on pH, ionic strength, and the presence of divalent cations. Examples of lipids useful in liposome production include phosphatidyl compounds, such as phosphatidylglycerol, phosphatidylcholine, phosphatidylserine, phosphatidylethanolamine, sphingolipids, cerebroside, and gangliosides. Particularly useful are diacylphosphatidylglycerols, where the lipid moiety contains from 14-18 carbon atoms, particularly from 16-18 carbon atoms, and is saturated. Illustrative phospholipids include egg phosphatidylcholine, dipalmitoylphosphatidylcholine and distearoylphosphatidylcholine.

An immunogenic target may also be administered in combination with one or more adjuvants to boost the immune response. Exemplary adjuvants are shown in Table I below:

Table I*Types of Immunologic Adjuvants*

Type of Adjuvant	General Examples	Specific Examples/References
1 Gel-type	Aluminum hydroxide/phosphate ("alum adjuvants")	(Aggerbeck and Heron, 1995)
	Calcium phosphate	(Relyveld, 1986)
2 Microbial	Muramyl dipeptide (MDP)	(Chedid et al., 1986)
	Bacterial exotoxins	Cholera toxin (CT), <i>E.coli</i> labile toxin (LT)(Freytag and Clements, 1999)
	Endotoxin-based adjuvants	Monophosphoryl lipid A (MPL) (Ulrich and Myers, 1995)
	Other bacterial	CpG oligonucleotides (Corral and Petray, 2000), BCG sequences (Krieg, et al. <i>Nature</i> , 374:576), tetanus toxoid (Rice, et al. <i>J. Immunol.</i> , 2001, 167: 1558-1565)
3 Particulate	Biodegradable polymer microspheres	(Gupta et al., 1998)
	Immunostimulatory complexes (ISCOMs)	(Morein and Bengtsson, 1999)
	Liposomes	(Wassef et al., 1994)
4 Oil-emulsion and surfactant-based adjuvants	Freund's incomplete adjuvant	(Jensen et al., 1998)
	Microfluidized emulsions	MF59 (Ott et al., 1995)
		SAF (Allison and Byars, 1992) (Allison, 1999)
	Saponins	QS-21 (Kensil, 1996)
5 Synthetic	Muramyl peptide derivatives	Murabutide (Lederer, 1986) Threony-MDP (Allison, 1997)
	Nonionic block copolymers	L121 (Allison, 1999)
	Polyphosphazene (PCPP)	(Payne et al., 1995)
	Synthetic polynucleotides	Poly A:U, Poly I:C (Johnson, 1994)

5 The immunogenic targets of the present invention may also be used to generate antibodies for use in screening assays or for immunotherapy. Other uses would be apparent to one of skill in the art. The term "antibody" includes antibody fragments, as are known in the art, including Fab, Fab₂, single chain antibodies (Fv for example), humanized antibodies, chimeric antibodies, human antibodies, produced by several methods as are known in the art.

10 Methods of preparing and utilizing various types of antibodies are well-known to those of skill in the art and would be suitable in practicing the present invention (see, for example, Harlow, et al. *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988; Harlow, et al. *Using Antibodies: A Laboratory Manual, Portable Protocol No. 1*, 1998; Kohler and Milstein, *Nature*, 256:495 (1975)); Jones et al. *Nature*, 321:522-525 (1986);

Riechmann et al. Nature, 332:323-329 (1988); Presta (Curr. Op. Struct. Biol., 2:593-596 (1992); Verhoeyen et al. (Science, 239:1534-1536 (1988); Hoogenboom et al., J. Mol. Biol., 227:381 (1991); Marks et al., J. Mol. Biol., 222:581 (1991); Cole et al., Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, p. 77 (1985); Boerner et al., J. Immunol., 147(1):86-95 (1991); Marks et al., Bio/Technology 10, 779-783 (1992); Lonberg et al., Nature 368 856-859 (1994); Morrison, Nature 368 812-13 (1994); Fishwild et al., Nature Biotechnology 14, 845-51 (1996); Neuberger, Nature Biotechnology 14, 826 (1996); Lonberg and Huszar, Intern. Rev. Immunol. 13 65-93 (1995); as well as U.S. Pat. Nos. 4,816,567; 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; and, 5,661,016). The antibodies or derivatives therefrom may also be conjugated to therapeutic moieties such as cytotoxic drugs or toxins, or active fragments thereof such as diphtheria A chain, exotoxin A chain, ricin A chain, abrin A chain, curcin, crotin, phenomycin, enomycin, among others. Cytotoxic agents may also include radiochemicals. Antibodies and their derivatives may be incorporated into compositions of the invention for use *in vitro* or *in vivo*.

Nucleic acids, proteins, or derivatives thereof representing an immunogenic target may be used in assays to determine the presence of a disease state in a patient, to predict prognosis, or to determine the effectiveness of a chemotherapeutic or other treatment regimen. Expression profiles, performed as is known in the art, may be used to determine the relative level of expression of the immunogenic target. The level of expression may then be correlated with base levels to determine whether a particular disease is present within the patient, the patient's prognosis, or whether a particular treatment regimen is effective. For example, if the patient is being treated with a particular chemotherapeutic regimen, an decreased level of expression of an immunogenic target in the patient's tissues (i.e., in peripheral blood) may indicate the regimen is decreasing the cancer load in that host. Similarly, if the level of expression is increasing, another therapeutic modality may need to be utilized. In one embodiment, nucleic acid probes corresponding to a nucleic acid encoding an immunogenic target may be attached to a biochip, as is known in the art, for the detection and quantification of expression in the host.

It is also possible to use nucleic acids, proteins, derivatives therefrom, or antibodies thereto as reagents in drug screening assays. The reagents may be used to ascertain the effect of a drug candidate on the expression of the immunogenic target in a cell line, or a cell or tissue of a patient. The expression profiling technique may be combined with high throughput screening techniques to allow rapid identification of useful compounds and

monitor the effectiveness of treatment with a drug candidate (see, for example, Zlokarnik, et al., Science 279, 84-8 (1998)). Drug candidates may be chemical compounds, nucleic acids, proteins, antibodies, or derivatives therefrom, whether naturally occurring or synthetically derived. Drug candidates thus identified may be utilized, among other uses, as pharmaceutical compositions for administration to patients or for use in further screening assays.

Administration of a composition of the present invention to a host may be accomplished using any of a variety of techniques known to those of skill in the art. The composition(s) may be processed in accordance with conventional methods of pharmacy to produce medicinal agents for administration to patients, including humans and other mammals (i.e., a "pharmaceutical composition"). The pharmaceutical composition is preferably made in the form of a dosage unit containing a given amount of DNA, viral vector particles, polypeptide or peptide, for example. A suitable daily dose for a human or other mammal may vary widely depending on the condition of the patient and other factors, but, once again, can be determined using routine methods.

The pharmaceutical composition may be administered orally, parentally, by inhalation spray, rectally, intranodally, or topically in dosage unit formulations containing conventional pharmaceutically acceptable carriers, adjuvants, and vehicles. The term "pharmaceutically acceptable carrier" or "physiologically acceptable carrier" as used herein refers to one or more formulation materials suitable for accomplishing or enhancing the delivery of a nucleic acid, polypeptide, or peptide as a pharmaceutical composition. A "pharmaceutical composition" is a composition comprising a therapeutically effective amount of a nucleic acid or polypeptide. The terms "effective amount" and "therapeutically effective amount" each refer to the amount of a nucleic acid or polypeptide used to induce or enhance an effective immune response. It is preferred that compositions of the present invention provide for the induction or enhancement of an anti-tumor immune response in a host which protects the host from the development of a tumor and / or allows the host to eliminate an existing tumor from the body.

For oral administration, the pharmaceutical composition may be of any of several forms including, for example, a capsule, a tablet, a suspension, or liquid, among others. Liquids may be administered by injection as a composition with suitable carriers including saline, dextrose, or water. The term parenteral as used herein includes subcutaneous, intravenous, intramuscular, intrasternal, infusion, or intraperitoneal administration.

Suppositories for rectal administration of the drug can be prepared by mixing the drug with a suitable non-irritating excipient such as cocoa butter and polyethylene glycols that are solid at ordinary temperatures but liquid at the rectal temperature.

The dosage regimen for immunizing a host or otherwise treating a disorder or a disease with a composition of this invention is based on a variety of factors, including the type of disease, the age, weight, sex, medical condition of the patient, the severity of the condition, the route of administration, and the particular compound employed. For example, a poxviral vector may be administered as a composition comprising 1×10^6 infectious particles per dose. Thus, the dosage regimen may vary widely, but can be determined routinely using standard methods.

A prime-boost regimen may also be utilized (WO 01/30382 A1) in which the targeted immunogen is initially administered in a priming step in one form followed by a boosting step in which the targeted immunogen is administered in another form. The form of the targeted immunogen in the priming and boosting steps are different. For instance, if the priming step utilized a nucleic acid, the boost may be administered as a peptide. Similarly, where a priming step utilized one type of recombinant virus (i.e., ALVAC), the boost step may utilize another type of virus (i.e., NYVAC). This prime-boost method of administration has been shown to induce strong immunological responses.

While the compositions of the invention can be administered as the sole active pharmaceutical agent, they can also be used in combination with one or more other compositions or agents (i.e., other immunogenic targets, co-stimulatory molecules, adjuvants). When administered as a combination, the individual components can be formulated as separate compositions administered at the same time or different times, or the components can be combined as a single composition.

Injectable preparations, such as sterile injectable aqueous or oleaginous suspensions, may be formulated according to known methods using suitable dispersing or wetting agents and suspending agents. The injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent. Suitable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution, among others. For instance, a viral vector such as a poxvirus may be prepared in 0.4% NaCl. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed, including

synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

For topical administration, a suitable topical dose of a composition may be administered one to four, and preferably two or three times daily. The dose may also be administered with intervening days during which no dose is applied. Suitable compositions may comprise from 0.001% to 10% w/w, for example, from 1% to 2% by weight of the formulation, although it may comprise as much as 10% w/w, but preferably not more than 5% w/w, and more preferably from 0.1% to 1% of the formulation. Formulations suitable for topical administration include liquid or semi-liquid preparations suitable for penetration through the skin (*e.g.*, liniments, lotions, ointments, creams, or pastes) and drops suitable for administration to the eye, ear, or nose.

The pharmaceutical compositions may also be prepared in a solid form (including granules, powders or suppositories). The pharmaceutical compositions may be subjected to conventional pharmaceutical operations such as sterilization and/or may contain conventional adjuvants, such as preservatives, stabilizers, wetting agents, emulsifiers, buffers etc. Solid dosage forms for oral administration may include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound may be admixed with at least one inert diluent such as sucrose, lactose, or starch. Such dosage forms may also comprise, as in normal practice, additional substances other than inert diluents, *e.g.*, lubricating agents such as magnesium stearate. In the case of capsules, tablets, and pills, the dosage forms may also comprise buffering agents. Tablets and pills can additionally be prepared with enteric coatings. Liquid dosage forms for oral administration may include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs containing inert diluents commonly used in the art, such as water. Such compositions may also comprise adjuvants, such as wetting, sweetening, flavoring, and perfuming agents.

Pharmaceutical compositions comprising a nucleic acid or polypeptide of the present invention may take any of several forms and may be administered by any of several routes. In preferred embodiments, the compositions are administered via a parenteral route (intradermal, intramuscular or subcutaneous) to induce an immune response in the host. Alternatively, the composition may be administered directly into a lymph node (intranodal) or tumor mass (*i.e.*, intratumoral administration). For example, the dose could be administered subcutaneously at days 0, 7, and 14. Suitable methods for immunization using compositions comprising TAs are known in the art, as shown for p53 (Hollstein et al., 1991),

p21-ras (Almoguera et al., 1988), HER-2 (Fendly et al., 1990), the melanoma-associated antigens (MAGE-1; MAGE-2) (van der Bruggen et al., 1991), p97 (Hu et al., 1988), and carcinoembryonic antigen (CEA) (Kantor et al., 1993; Fishbein et al., 1992; Kaufman et al., 1991), among others.

5 Preferred embodiments of administratable compositions include, for example, nucleic acids or polypeptides in liquid preparations such as suspensions, syrups, or elixirs. Preferred injectable preparations include, for example, nucleic acids or polypeptides suitable for parental, subcutaneous, intradermal, intramuscular or intravenous administration such as sterile suspensions or emulsions. For example, a recombinant poxvirus may be in admixture
10 with a suitable carrier, diluent, or excipient such as sterile water, physiological saline, glucose or the like. The composition may also be provided in lyophilized form for reconstituting, for instance, in isotonic aqueous, saline buffer. In addition, the compositions can be co-administered or sequentially administered with other antineoplastic, anti-tumor or anti-cancer agents and/or with agents which reduce or alleviate ill effects of antineoplastic, anti-tumor or
15 anti-cancer agents.

A kit comprising a composition of the present invention is also provided. The kit can include a separate container containing a suitable carrier, diluent or excipient. The kit can also include an additional anti-cancer, anti-tumor or antineoplastic agent and/or an agent that reduces or alleviates ill effects of antineoplastic, anti-tumor or anti-cancer agents for co- or
20 sequential-administration. Additionally, the kit can include instructions for mixing or combining ingredients and/or administration.

A better understanding of the present invention and of its many advantages will be had from the following examples, given by way of illustration.

EXAMPLES

Example 1

A. Modification of mCEA (6D) repeat 1

The presence of truncated forms of CEA in cells following expression of recombinant
30 CEA has been documented. This study set forth to generate CEA-encoding nucleic acid sequences that do not result in the expression of truncated CEA following expression in cells. Generation and expression of a new CEA-encoding nucleic acid sequence, CAP(6D)-1,2 (SEQ ID NO:28), is described below.

The plasmid p3'H6MCEA was obtained from Virogenetics, Inc. This plasmid contains the MCEA gene with 6D modification under the control of partial H6 promoter (**Fig. 1A**; SEQ ID NO: 23). The 912 bp NruI – BamHI fragment from p3'H6MCEA was cloned into pUC18 to form plasmid pSE1544.9 (pUC18-mCEA repeat 1; **Fig. 1B**).

5 OPC purified Oligos 7524-7526, 7528-7533, 7535-7537, and 7567-7568 were kinased and annealed to create two fragments which were ligated to result in a 464 bp synthetic modified mCEA repeat 1 flanked by AccI and BamHI sites. This synthetic modified repeat 1 fragment was cloned into pSE1544.9 AccI-BamHI to create pSE1616.44 (pUC18-mCEA-modified repeat 1; **Fig. 2**). The 904 bp EcoRV – BamHI fragment of pSE1616.44 was cloned
10 back into p3'H6MCEA EcoRV-BamHI to form pSE1658.15 (p3'H6MCEA-modified repeat 1; **Fig. 3**).

15 **B. Modification of mCEA(6D) repeat 2**

A synthetic modified repeat 2 fragment was created by using a method called gene splicing by overlap extension (SOE) and cloned into pBluescript-SK+, generating pBSmCEA (**Fig. 4**). The oligos used for the repeat 2 modification are shown below (section IV, B). The two different clones (pBS-mCEA-3 and pBS-mCEA-8) contained various point mutations.

20 The 697 bp BamHI – EcoRI fragment of pBS-mCEA-3 was cloned into pUC18 BamHI – EcoRI to create pSE1671.8. The 591 bp SpeI – Bsu36I fragment of pBS mCEA-8 was cloned into pSE1671.8 SpeI-Bsu36I, generating plasmid designated pSE1681.1. Two site PCR mutagenesis, using the Quikchange site directed mutagenesis kit from Stratagene with oligos
7751

25 (GGACGGTAGTAGGTGTATGATGGAGATATAGTTGGGTCGTCTGGGCC; SEQ ID NO: 24) and 7760 (CAGAATGAATTATCCGTTGATCACTCC; SEQ ID NO: 25), was performed to correct the two remaining point mutations pSE1681.1. The corrected clone was designated pSE1686.1 (pUC18 mCEA modified repeat 2; **Fig. 5**).

30 As noted recently, an Alanine codon was absent from 5' terminus of the second repeat in plasmid p3'H6MCEA which contained CEA. To preserve the consistency of the amino acid sequence of CEA, the Alanine codon present in plasmid pSE1686.1 containing the modified second repeat of CEA was knocked out. This was accomplished using oligos 7802 (CGTGACGACGATTACCGTGTATGAGCCACCAAAACCATTTCATAAC; SEQ ID NO:

26) and 7803 (GTTATGAATGGTTTTGGTGGCTCATACACGGTAATCGTCGTCACG; SEQ ID NO: 27) and the Quikchange site-directed mutagenesis kit from Stratagene. The resulting plasmid, pSE1696.1 (pUC18 mCEA modified repeat 2; Fig. 6) was confirmed by sequencing.

5 The 694 bp Bsu361-BamHI fragment from pSE1696.1 was cloned into Bsu361-BamHI site of Pse1658.15 to combine modified repeats 1 and 2. The generated plasmid was designated p3'H6modMCEA-1st&2nd repeats (Fig. 7).

C. Construction of ALVAC donor plasmid pNVQH6MCEA(6D1st&2nd)

10 The 2.2 kb NruI/XhoI fragment from p3'H6modMCEA-1st&2ndrepeats was cloned into NruI/XhoI site of pNVQH6LSP-18, generating pNVQH6MCEA(6D1st&2nd; Fig. 8). The modified CEA sequence ("CAP(6D)-1,2") contained within pNVQH6MCEA is shown in Fig. 9 and SEQ ID NO:28.

D. Expression of modified CEA

15 To test the stability of the CAP(6D)-1,2 sequence upon expression in a cell, the gene together with flanking H6 promoter was PCR amplified using pNVQH6MCEA(6D1ST&2ND) as template and two oligos (8034LZ: CTGGCGCGCCTTCTTTATTCTATACTTAAAAAGTG (SEQ ID NO: 29); and 8035LZ: CTGGTACCAGAAAACTATATCAGAGCAACCCCAAC (SEQ ID NO: 30)). The PCR product was then cloned into an NYVAC TK donor plasmid designated pLZTK1 containing the LacZ and K1L marker genes. This vector was specifically made for the generation of recombinant virus in NYVAC by using blue/white screening method. After in vitro recombination between donor plasmid pLZTK1mCEA(6D1st&2nd) and NYVAC, the foreign CAP(6D)-1,2 sequence and marker genes are integrated into the NYVAC genome. The plaques containing intermediate recombinant NYVAC with both LacZ and mCEA appeared blue. Several rounds of plaque purification were then performed. The second recombination event kicked out the marker genes resulting in the final white plaques containing recombinants with only the CAP(6D)-1,2 sequence but no marker genes (Fig. 10).

30 The recombinant white plaques and blue plaques were picked for confirmation of CAP(6D)-1,2 sequence expression. Infection was performed using the virus from the respective plaques and the cells were harvested three days after infection for preparing either

cellular DNA or cell lysate. For isolation of recombinant NYVAC DNA, DNAzol® reagent (GibcoBRL) was used. PCR (PCR Condition: 95°C (5min)→ [95°C(30sec) → 49°C(30sec) → 72°C(1min)] 30 cycles→72°C (7min)→ 4°C) was run to confirm the existence of CAP(6D)-1,2 sequence in the recombinant NYVAC genome. The primers used were 7569LZ (5' ttggatccatggaggtctccctcgccc 3' forward primer; SEQ ID NO: 31) and 7570LZ (5' ttggatccctatatcagagcaacccc 3' reverse primer; SEQ ID NO: 32), which could amplify the full length 2106 bp CAP(6D)-1,2.

Comment [AW1]:

The final recombinant white plaques PRBC-III- 2, 3, 6, 8, 9, 10 all demonstrated the 2.1 kb CAP(6D)-1,2 sequence band in PCR. PRBC-III-N1 was a blue plaque with both marker genes and CAP(6D)-1,2 sequence still in the viral genome and the CAP(6D)-1,2 sequence band was also amplified in the PCR. The prominent PCR band amplified from vCP 307 DNA (containing native CEA integrated into the ALVAC genome) was truncated CEA at 1.2 kb with a very faint full-length CEA band. The cell-only sample (no viral infection) was used as a negative control and the plasmid pLZTK1MCEA(6D1ST&2ND) was a positive control used in the PCR reaction. The PCR results clearly showed the full-length CAP(6D)-1,2 in the recombinant viral genome with no other truncated form of CEA visible. This result indicated that CAP(6D)-1,2 has increased stability relative to the native CEA in the ALVAC genome.

Protein expression was also assayed by immunoblot to confirm the absence of truncated CEA protein in cells expressing CAP(6D)-1,2 (Fig. 11). For isolation of cell lysate, cells were first washed with PBS followed by the addition of Lysis Buffer (Reporter Gene Assay; Boehringer Mannheim) and shaking for 15 minutes. Cell lysate was spun down at 13,000 rpm and the supernatant was collected for Western blot analysis. Samples were loaded onto a 10% polyacrylamide gel and run at 125 volts. The protein was then transferred to a PVDF filter membrane (Immobilon-P, Millipore). An HRP-linked mouse CEA monoclonal antibody (1:1000; Fitzgerald) was used to detect the expression of mCEA with the enhancement from a chemiluminescence reagent (DNA Thunder™; NEN™ Life Science Products).

All six final CAP(6D)-1,2 recombinant white plaques (PRBC-III-2,3,6,8,9,10) and one intermediate blue plaque (pRBC-III-N1) showed only one CEA band with no other truncated form (Fig. 11). In contrast, protein from vCP307 plaques (recombinant ALVAC expressing native CEA) showed a clear truncated CEA product at ~60 kDa in addition to the

full length CEA. Prolonged exposure of the film verified the absence of any truncated CEA polypeptides in the CAP(6D)-1,2 recombinants. CEF was used as the negative control.

5 In conclusion, the CAP(6D)-1,2 recombinants were generated with the mCEA instead of the native CEA to prevent the expression of multiple versions of CEA. CAP(6D)-1,2 expressed from recombinant NYVAC was proven effective in eliminating truncated version of CEA by both PCR and Western blot.

10 While the present invention has been described in terms of the preferred embodiments, it is understood that variations and modifications will occur to those skilled in the art. Therefore, it is intended that the appended claims cover all such equivalent variations that come within the scope of the invention as claimed.

~~SEQUENCE LISTING~~

~~SEQ ID NO.: 1, AAC2-1 nucleotide sequence~~

Seq ID: R011_17_A02_1_Nucleotide_Sequence
AGCAGGACGGGGGGCTGTGTGGCTATCGGCTTCGGCGGGGGGGGGGGAGGGAGCGGCTGGGGTACGTTGGCGGAGTTG

5 AG
TGGGCACTCCTCGGACGTTGCTGGGCAACCTGAACGAGCTGGGGCTGGGGGGGATCGTCACTCAAGTCAAGCTGCT
GG
TTGGGGGGCAACCGCTGACAGCAGACAAAGGACTTGTATCGGCTGGAGTGGCTTCTTGTATTCAATTTTGGGG
GG
CGTGGGGGAGTGGGGCTGGACGTTGCTGTCTGTGGGGGGGGTGGGAAAGGAGAGGGTTGGGGGGTGTATTGGAG
TT
GATGTAGAGTTGGGGGCTGGGGCTGTCTGTGAGGCACTGCACGAGGAGTCTAGCGGGGGGGACCTATTTGGACAT
GG
AGGACGTCGTCGAGCGATGCCACGGCTTCATGCAGGGCAGCTATCAAGCTCTGGGATCTCGCTGGGGGGGCTGG

15 AA
GCAGAACGGGGCAACAGGGCAACGGGGGGCTGCAGCAGGTAGTCCGAGGGGCTGGGAAAGGAGACCCAGACCCAGCT
AG
TGAATCTGGAAGTGCAGTCAAGGGGGGGGAGTCCAGGCAGGGCTGACGGCAAGGGCTGCAACTGCAAAAAGTA
GA
ACTACATGCTGCTAAAGTCTCAGGGCTGCCAAGCAGGAGCGTCTGTGGGAGAGAACTTCTGCTCAACCTTGGG
GG
CAAGCCAGGCTCGGGAGTGGAGACGAGGGCTCCAGCAGCAGGAGCAGCAGCAGCAGCAGCAGCAGTCAAGAGGA
GG
CATTCCTGCTGGGAGAGCAGGCTCTCTCCAACCTGCTGCCACTCTGCACTTCAATGTGGGGCTGGAGCCAGTAC

25 GG
CCTACCTGCTGAGATCCGAGGCTCAAGACACCTCTGGATCAGGCTCTGAAGGGGCTGCTGCACTAGCGGGAGTGA
AT
TTTTACCTGCCAGAACTGTGAGGCTGTGGCAGGCTGCTCATCGGGGGCTGGAGTCTTGGTTCTTCTGGGAGCAA
GA
CAAACCTTATAAGTGTGACCTGTGGCGGTCTTGCTTGGGCTAGAACGGCAAGCTTGGCAGTCACCGTACAGTCCA
GA
CAGGGCAAAAGCGTTACCACTGCTCAATCTGGCGAGCGGCTTTTAACGGGGCAGCAAACTGAAAAGCCAGAGCG
GG
ATCCATTGGGAGAGAAAGCGCTATAAGTCTGAGACCTGGGGCTGGGGCTTTGTAGAGCTGGCAGATCTGGGGGG
35 GA
CGTCTGATCCAGACCGGGGAGAAAGCGCTACCGTTGGGCTAGCTGGGGAACCGGCTTCCGGCACCTGCAGAGCGT
GA
AGAGGCAGCTTGGCATGAGACAGCGGAGAGAAAGGCTTAGCACTGGCAGCGGCTGTGGGCTGCATTTCGGGCAGAA
GT
CAAGTGGGGCTGGATCTGGGGCAGAAACACGGAGCTGCTAGCAACAGCAAACTGCATACCAATTTCTGGGGGG
40 GG
GTAGGTCAGCGGAGGCGGACGGCGGCACTTGGTTCTTGGGCTGGGAAAGCTGCAGGGGAGGGCTTGGTTGGCTA
TG
AGGCTTGGGATAGCGGCTGTGGAGCGGCACTTTGCTATCAGAAATTGCCACCGCTCTTAATTTCTGACTGGGACA
45 GG
AGCGGTGGGAGATGCTGGCTACATCTGGCTGTCTTTTGGTGGTCANAGCGCTTTCGGCAGAACGGAGATTGTTTC
TG
AGGAGAGAGCTAGCTAGGGGCTGGCAAAAGGGGAGAGATTTCAGTCTGCTGCTTAAAGGAATAGCGCTCGAGG
TG
TGGGGGGCATTCGATTGAGTTTATCTGTAAATATAATTTATTGAGGGCTTTGGGTCGCACGGGGGGCTTCATTG
50 GA
TTGGATTTCCCACTCCGCTCTTCCACAAGTCTGATTAAAACTGACCAGAAAGCAGAGAGCTGAGATCAGAGCTCT
GG
TGGGAGAGATTACTAGCGGCTGGGCTCTCTGCTTTGGCTTGGCTATTTTATATTATTCTCTGATAACTTTTATCT
55 TT
AGAATTTCTGTTTCTCTGCTTTCTTTGGCTTCTTACTTTGTTTAAANTGCAAAAAGGGGTTCTCTGCTTCTGCGG
CT
CTAATTTCTAGCTCTGCAAGCTTTATTTCTTCTAGGGCAGCTCTGGCAACATCGGGGATTCGCAATTCGGCTCAGG
AA

CCGTCTCTGCTATTCTGGATGTTCTAGGTTCTCTACCACTCTAGAAATCGATACACACATTTCTCTCTTCTTCAA
 GG
 CTGATACGAACCATTTATGTTGACGGCAAAATCGAAGTAATAATAATCGCTGCTGCACGGCTCTCGGCTCTGGCGCA
 TT
 5 CTGTATCTGGATTGCGGTATCACTCGAACTCGACGGCTGTGGGTCTGGCGGATTGCTGTATCTGCATTCCGTATCACT
 GG
 AACTCGACGGCTCGGACGTTTCTCTGCAACATCGTGCAGAACTCTAAATCTCCGATTAATCTCCTCACTTGGCTTT
 GG
 CTCTGCTGTATCCATCTATAGTCTAGACAGCCAGCAGCGCTCAACTCGACTGCATCATCTCCACGGCGGGCT
 10 GT
 TCTTAGTACTGAGTTGATCGCTCGATCGGGCAGACATCAGACATTGCTTATCAGACATCATCTCAGCTTTTCTGA
 CT
 CTGGCCACTCTCTATGAATGTTATGGCCTAGCGCAAGAATCATCAAACTCTTTAGCTTCATTAGATCGTAAACAGT
 CT
 15 TAACGCATCGTTTACTACAGACCCATATCGGCTTTGAATCTTACCTGGCGTTCTCTGTATTGACTTGAACCGGCTTC
 TT
 CCTTTAGTGGCTTTTGGACATCTTCTGCGCACTCTCCACATGCCAGAACCTTCTTTTCTCTAGAACGGCATGCTG
 CT
 TCGTAACGTTAGCTTTTAAACGCTGGGTCTCTCAGCTGCTCTTGGCATCGCTGCATTGCTCTCTGGAACGAGTGA
 20 AT
 GCATTAGAACCTTGCATAGCAAAAGAAAAGGGGCTGAGTTCCATTCTCGGTTTGGTCTAGTTTGGTTGGCATTAT
 TG
 TTGGCATTACACATCTAAAGATTCACTAGCGCCATAGCCCAAGCGCTGTTCTAGTTGACCAACTTTCAAGTAGG
 AT
 25 TAAGAGGTTGGTTGACGGCTGCACTTTCTGCTCTAGCCGAGGTAGCTAGAAAGTCAGGAACACGGTTGGCTCTTG
 GG
 TGGCTGGAGTCTCTCAAACTTTAGAACGAAGCGCTCAAGCGTTGATTGATAGTTCTGGCGGTTGTTGGGGTGGGGG
 TT
 ATCTGATTATGGGACGACGCTAGAAACTAAGAAGCACTTTTGAATTTCTGGGCTAGAACTTCAACAATAAGTCAG
 30 TT
 CTAGTGGCTCTCGGCTCGGGACTAGTCAGAAAGCTACTCTTCTCGCTCTTGGCTCTTTCTGGCCATGGGGGCACT
 GG
 ACAATTAAAGAACGAAGCAAGCGAAGGGCGAGGACTCTATAAGAACGAATCATGATTCTATTTAGCAGATTGGAT
 GG
 35 GCAGCTGCAGAAATCGCTCGGGCTAGAAATCTTACATCTTGCAGCATCAGATCCTTGCATAAAGAACGCTCTCTG
 GG
 CAAAAAAAAAAAAAAAAAAAA

ATGGGTTTGGGGGGGGGGGGGAGGACGGCTGGGCTACGTTGGGCACTTCACTTGGGCACTGCTGGGAGGTGGTGGGCAAGGCT
CAACGAGCTGGGGCTGGGGGGGATCTCTCACTACGCTACGGCTGCTGGTGGTGGGGGGCAACGGCTCAGACGACAGAGGCACTTC
TCATGGCTCGACATGGCTTCTTATTACATTTTGGGGGGGCTGGGCACTCGGGCTACAGCTCTCTCTCTGGGGGGGGCT
GGGCAAGGCAAGGCTTGGGGGGCTCTATTGGACTTCATGTAACATTTGGGGGCTGGGGCTCTCTCGACGGCACTGCACAGGAC
CTACGGCGGGGGGGGCTATTTCGACATTCGACAGGCTGCTCCAGGCATCCGACGGCTTCATTCAGCGGCACTTCAACGCTTCG
GCATTCCTGGTGGGGGGCTTCGACAGCAAGCGGCAAGCGGCAAGCGGCTTCACAGGCTACTGGGAGGGGCTGGGACGA
CAGCCAGACCCACCTACTCAATCTCGAAGCTCGACTCAAGCGGGGGGGCACTTCGACGCACCGCTGACCCGAAGGGCTGGAACTG
CAAAAGCTACAACTACATGCTGCTAAACTCTCAGGGCTGCCAAGCAGCGGAGCGTGGTGGGGGGCAAGCACTTCTCGCTCAAGCTT
GGGGCAAGCAAGCTCGGCACTCGAGACGCTCGACAGGACGAGCAGCAGCAGGACGAGCACTGAAGAGAGAGGCAAT
GCTGCTGGGCAAGGAGGCTCTCTCGAACTGCTGGCACTCTCGACTTCAAATCTGGGGGCTCGAGGCACTTACGGGCTAGCTGCT
GACTTCGCAAGGCTCAACAGACCTCTCGGATCAGGCTCTGCTACGGGGCTGCTGGCACTACGGGCAAGTCAATTTTTGAGGTGGGCA
ACATTCAGGCTCTGGGAGGCTGCTATCGGGGGCTGCACTGCTGCTGGGCAAGCAAGCAAAAGCTATAACTCTPACGCTG
TGGGGCTCTGCTTGGGCTACAGGGCAAGCTTGGGACTCATGCTACAGTCGCAAGCGGGGAAAGGCTTACAGACTGCTCAAT
CTGGCGAGGGGCTTTTAAAGCGGGCAAGCAAGCTGAAAGCGCAAGCGGCTTCATTTGGGCAAGAGCGGCTATAAGCTGAG
GCTGGGCTGGGGCTTTCTACAGCTGGCAATCTCGGGGGCAAGCTGCTGATCAGACGGGGCAAGGCTTACCTTGGGCT
ACCTGGGCAAGCGGCTTGGGCAAGCTCGACAGGCTCAAGAGGCAAGCTTGGCATCGACAGCGGAGAGAGGCTTACCACTGGGA
GGGCTCTGGGCTGGATTTCGGGCAAGCAAGCTCAAGTGGGGCTGCATCTGGGGCAAGAACAGCGAGCTGCTACGACAGCAAG
TGCAGTACGACATCTCTGGGGGGGGGCTAG

~~ATACCGCGAACTCCCTAAGCCTTCTATTAGCTGCAATAATAGTAAGCCTGTCCAAGACAAAGATG~~

CCCTGTGTCCCTAGACTCGAAGTCAAGCAACCGAAATAGAACTCTCACCGTCTTTAACCTCAGCAGGAAG

~~ACGTGCTTTACGGACCCGATCGTGCTACAATCAGGCCTCTAAACACAAGCTATAGATCAGCGGAAAATCT~~

~~ACGTTAAACAGGGTCAGAGTTCTATTTCCGTTGCTGAGTTGGAGTCTAGCGGACACAGGCAGGCAGTGGT~~

CTCATCTATAGCTTGTGTTTAGAGCGCTGATTGTAGGAGCATCCGGTCCGTAAAGCAGCTTCAGAATCAG

~~GATCCACTATTCTTCACCGTAATATTCCGAATCAACACTTCCTGGGTGGACTGTTGGAAACTG~~

SEQ ID NO: 37 (353)
GACACAGCAAGCTAGAAATGCCAAACCCAAAATCCAGTCAGCGCCAGCAGGTCTGATTTCAGTCATTCTCA

~~TGAATCAGACCTCCTGGCGCTGACTCGATTTTGGGTTTCGCATTTCTAGCTTCCTGCTCGGTTCCCTGGTC~~

GATGCTACACCTGCCAAGCTCACAATACGGACACCGGACTGAACCGGACAACCGTGAAGGACGATTACCGTGTATG
 CCGA

~~CATCCTCAACTCGGTTAGAATTGTTACTACTTATGAATCGTTTTGCTGGCTCGGCATACACGGTAATCGT~~

CTGGG

CTGTATATCATAACCCGACATTGACACTCGCTGTCGGCTTACTCGCAATCATGTAGCACCTTATCACTCTGGCATTCA
GAATG

~~SEQ ID NO.: 15, 7679~~

~~TTTGTATGGCCAGACGACCAACTATATCTGCATCATACCTACTACGGTCCGGGGCTGAAGTTGAGCGTTTC
TTGGG~~

5 ~~SEQ ID NO.: 16, 7680~~

~~TCATGCCAATTCACCAGCATACTCAAGACTTATTTATAAGCAACATAACTCAGACGAACACCGGACTCTATAG
TTGGG~~

~~SEQ ID NO.: 17, 7681~~

10 ~~TAAACAAATAACTGTTTCGGCCGAGCTGCCCAACGGCTCCATCTCCAGCAACAACGCCAAGCCGTCGAGCAGAA
GCATG~~

~~SEQ ID NO.: 18, 7682~~

15 ~~ATGTGGGTTATCATATAGAGAACTGCAAGCGCTCGGCTAACCGCCAAACTTTGCTTATTGAGCCAGCATAAATAAG
TCGTA~~

~~SEQ ID NO.: 19, 7683~~

20 ~~GCTCGTCTGGGCCATACAAACATTAACGATAAGAGCGCTGGGACTGATCAAGGCATAATTGATTCTGAATGCCAG
ACTCA~~

~~SEQ ID NO.: 20, 7684~~

~~GCTCGTCAATGTTTTCATCAATCAGCCAGCAGTACTGTCCACGGCGGTTGGATCGTGCATGCCAAGAAAGGCTCA
ACTTG~~

25 ~~SEQ ID NO.: 21, 7685~~

~~CGGAAGAGTTATTGTTTAAAGTCTAGTCTGCTGCTGACCACTCGCTGACTTATTGGGCTGGCAAGTATAGACTG
CGCTG~~

~~SEQ ID NO.: 22, 7686~~

30 ~~GCTCAGCTTCACAGCTGAAGGGCAGACCATCCTTCTGCTCGACGGGT~~

~~SEQ ID NO.: 23, CEA CAP6D~~

35 ~~ATGCACTCTC CCTCGGGCCG TCGGCACAGA TGGTGCATCC CCTGCCAGAG GCTCCTGGTG
AGACCGCTCAG TTCTAAGCTT CTGGAACCGG CCGACCACTG CGAAGCTCAG TATTGAATCG
ACGCGGTTCA ATGTGGCAGA GGGGAAGGAG CTGCTTCTAC TTGTCCACAA TCTGCGGCAG
CATCTTTTTC CTTACAGCTG CTACAAAGCT CAAAGACTGG ATGCCAAGCG TCAAATTATA
GCATATCTAA TAGGAAGTCA ACAAGCTAGC CCAGCGCCCG CATACAGTCC TCGACAGATA
ATATACCCCA ATGCATCCCT CTTGATCCAG AACATCATCC AGAATCAGAG AGGATTCTAG
ACGCTACAGG TCATAAAGTC AGATCTTCTG AATGAAGAAG CAAGTGGGCA GTTGGGGTA
40 ~~TACCGGGAGC TCGCCAAAGC CTGCATCTCG AGCAACAACCT CCAAGCGGCT GCAGCACAAG
GATGGTCTCG CTTTCACCTG TGAAGCTGAG ACTCAGGAGC CAACCTAGCT GTGGTGGGTA
AACAAATCAG CCGTCCCGCT CAGTCCCAGG CTGCAGCTCT CCAATGGCAA CAGCAGCCTC
ACTCTATTCA ATGTCAACAAG AAATCAGACA CGAAGCTACA AATGTGAAG GCAGAACGCA
CTCAGTCCCA GCGGCACTCA TTCAGTCATC CTGAATCTCC TCTATGGGGC GCATGCGGGC
45 ~~AGCATTTGCC CTCTAAACAG ATCTTACAGA TCAGCGGAAA ATGTCAACCT CTCCTGGCAG
GCAGCGTCTA ACCCAGCTCG ACAGTACTCT TGGTTTCTCA ATGGCACTTT CCAGCAATCG
AGCGAAGAGG TGTTTATCCG CAACATCACT CTGAATAATA CTGCATCGTA TACCTGGCAA
GCCCATAACT CAGACACTCG CTTGAAATAGG ACCAGACTCA CGAGCATCAG AGTCTATGAG
CGACCGAAAG CTTTCATCAG CAGCAAGAAG TCGAAGCCCG TCGAGCATCA CGATCGTGTA
50 ~~CCCTTAAGCT CTGAAGCTCA CATTGAGAA ACACCTAGC TGTGCTGGGT AATAATCAG
ACGCTCGGGC TCACTCCGAG CCTGCAGCTG TCGAATGAGA ACAGCAGGCT CAGTCTAGTC
ACTGTACAA CCAATCATCT AGCAGCCTAT CAGTCTCGAA TCGACAAGCA ATTAAGTCTT
GACCACAGCG ACCCAGTCTAT CTTGAATCTC CTCTATGGG CAGACGAGCG CAGCATTTCC
CGCTCATACA CCTATTACCG TCGAGGGCTG AACCTCAGCG TCTCCTGGCA TGCAGCGCTCT
55 ~~AACCCAGCTG CACACTATTG TTGGGTGATT GATGGGAAGA TCGACGAAGA CAGACAAGAG
CTGTTTATCT CCAACATCAG TCAGAGAAG AGCGGACTGT ATACCTGCCA GGCGAATAAG
TCAGCCAGTG CCCACAGCAG CACTACACTC AAGACAATCA CAGTCTCTCG CGAGCTGGCG
AAGCGCTCGA TCTCCAGCAA CAAGTCCAAA CCGGTCCAGC ACNAGCATCC TGTGGCGTTG
ACCTCTGAG CACAGGCTCA GAACAGAAGC TAGCTCTGCT CGCTAAATCG TCAGAGGCTG
60 ~~CCAGTCAGTC CCAGGCTGCA CCTGTCCAAT GGCACAGGA CCGTCACTCT ATTCAATCTG~~~~~~~~~~~~

~~ACAAGAAATG ACGGAACAGG CTATGTATGT CGAATCCAGA ACTCAGTCAG TCGAAACGGC~~
~~ACTCAGCCAG TCAGGCTGGA TGTCTCTAT CCGCGGAGCA CCGGCATCAT TTCCCCGGCA~~
~~GACTGCTGTT AGCTTTCGGG ACGCGAGGTC AAGCTGTGCT GGCAGTGGGC GTCTAACGGA~~
~~TCCCGGCACT ATTCTTGGGG TATCAATGGG ATAGCGGAGG AACACACAGA ACTTCTCTTT~~
~~ATCGCGAAAA TCACGGGAAA TAATAACGGG ACCTATCGGT CTTTCTCTGC TAACTTGGGT~~
~~ACTCGCGGCA ATAATTCCAT ACTCAACAGG ATCAGACTCT CTGCATCTGG AACTTCTCCT~~
~~GCTCTCTCAG CTGCGGGCAG TGTGGGCATC ATCATTCGAG TCCTGCTTGG CTTTCTCTGC ATATAG~~

SEQ ID NO.: 24, CAP6D 1,2

~~ATGGAGTCTG CCGCGGCGCG TCGGCACAGA TGCTGGATCG CCGCGGAGAG CCGCTGCTG~~
~~ACAGCGTCAG TTCTAACCTT CTGGACCGCG CCCAGCACTG CCAGCTCAG TATTGAATCG~~
~~ACGCGGTTCA CTGTGCGAGA CCGGAAGCAG GTGCTTCTAG TTCTGACAAA TCTGCGCGAG~~
~~CATCTTTTTC GCTACAGCTG CTAGAAAGGT GAAAGACTGG ATGCGAAGCG TCAAAATTATA~~
~~GCATATCTAA TAGGAAGTCA ACAGCTAGC CCAGCGCGCG CATACACTGG TCGACAGATA~~
~~ATATACCGGA ATGCATCGCT CCGTATCGAG AACATGATCG AGAATGACAG AGGATTCTAG~~
~~ACCGTACAGG TCATAAAGTC AGATCTTCTG AATCAAGAG CAACTGCGGA GTTCCGGGTA~~
~~TACCGGGAAC TCCCTAAGCG TTCTATTAGG TCGAATAATA TTACCGCTGT GGAACAGAAA~~
~~GATCGGGTGG CTTTTAGATG CGAGCGGCAA ACTCAAGAG CAAGATATCT CTGGTGGGTG~~
~~AACAACCACT CCGTCCCTGT CTCCGCTAGA CTCCAAGTCA GCAACGGAAA TAGAAGCTCG~~
~~ACCGTCTTTA ACCTGACCGA GAAGCAGACA CCAAGCTAGA AATGCGAAG CCAAAATGCA~~
~~CTCAGCGGCA CGAGCTCTGA TTCACTGATT CTCAGCTCG TTTACCGAGC CGATCCCTCT~~
~~AGAATGAGCG CTCTAAACAG AACCTATAGA TCAGCGGAAA ATCTGAATCT CAGCTCTCAT~~
~~GGCGCTAGCA ATCTGCGCGG CCAATACAGG TGCTTTGTGA ATCGGCACTT CCAACACTCG~~
~~ACCCAGCAAC TCTTCATTGG CAATATTAGG CTCAACAATA CTGGATCGTA CAGGTGCGAA~~
~~GCTCACAATA CGGACAGCGG ACTCAACCGG ACAAGCGTGA CGAGGATTAG CCGTATGAG~~
~~CGACCAAAAG CATTGATAAG TACTAACAA TCTAACCCAG TTGAGGATGA CGAGCGACTT~~
~~GCATTAACTT CTGAGCGAGA CATTGAAAAT ACGAGTTATT TATGGTGGGT CAATAACGAA~~
~~AGTTTGGCGG TTAGCGCAGG CTTGCACTTG TCTAATGATA ACCGCACATT GAGACTCCTG~~
~~TCCGTTACTG CGAATGATGT ACGACCTTAT CAGTCTCGCA TTCAGAATGA ATTATCCGTT~~
~~GATCACTCGG ACCCTGTTAT CTTAATCTT TTCTATCGCG CAGACGAGCG AACTATATCT~~
~~CGATCATAGA CCGTACCGCG TCGCGCGCTG AACTTGAGCG TTTCTTGGCA TCGACGATCG~~
~~AACCGCGCTG CACAGTACTG CTGGGTGATT GATGGAAACA TTCACCGCA TACTGAAGAG~~
~~TTATTTATTA GCAACATAAG TGAGAGAAAC ACGGACTCT ATACTTGGCA GCGCAATAG~~
~~TCAGCGACTG CTCACAGCAG CACTACACTT AAAACAATAA CTGTTTGGCG CGACCTGGCG~~
~~AAGCCCTGCA TGTCCAGCAA CAATGCAAAA CCGCTGGAGG ACAAGGATGC TCTGGCCTTG~~
~~ACCTGTGAAC CTCAGGCTCA CAACAGAACG TACCTGTGCT GCGTAAATGC TCAGACCGTC~~
~~CGAGTCAGTC CCAGGCTGCA CCTGTCCAAT GCGAACAGGA CCGTCACTCT ATTCAATGTC~~
~~AGAAGAAATG ACGGAACAGG CTATGTATGT CGAATCCAGA ACTCAGTCAG TCGAAACGGG~~
~~ACTCAGGCGA TCACCGTCCA TGTCTCTAT CCGCGGAGCA CCGGCATCAT TTCCCCGGCA~~
~~GACTCTGCTT ACCTTTCGGG ACGGAGCGTC AACCTCTGCT CCGACTCGCG GTCTAACGGA~~
~~TCCCCGCACT ATTCTTGGGG TATCAATGGG ATAGCGGAGG AACACACAGA ACTTCTCTTT~~
~~ATCGCGAAAA TCACCGGAAA TAATAACGGG ACCTATCGGT CTTTCTCTGC TAACTTGGGT~~
~~ACTCGCGGCA ATAATTCCAT ACTCAACAGG ATCAGACTCT CTGCATCTGG AACTTCTCCT~~
~~GCTCTCTCAG CTGCGGGCAG TGTGGGCATC ATCATTCGAG TCCTGCTTGG CTTTCTCTGC ATATAG~~

SEQUENCE LISTING

<110> Sanofi Pasteur, Ltd.
 Therion Biologics, Inc.

<120> Modified CEA Nucleic Acid and Expression Vectors

<130> API-01-20-US

<140> 10/510,677

<141> 2003-04-09
 <150> US 60/370,972
 <151> 2002-04-09
 5 <160> 32
 <170> PatentIn version 3.3
 10 <210> 1
 <211> 3564
 <212> DNA
 <213> Homo sapiens
 15 <220>
 <221> misc_feature
 <222> (1663)..(1663)
 <223> n is a, c, g, or t
 20 <400> 1
 agcaggaccg gggcctgtgt cgctatgggt tccccgccg ccccgaggag agcgctgggc 60
 tacgtccgcg agttcactcg ccaactcctcc gacgtgctgg gcaacctcaa cgagctgcgc 120
 25 ctgcgcggga tcctcactga cgtcacgctg ctggttggcg ggcaaccctc cagagcacac 180
 aaggcagttc tcatcgcttg cagtggcttc ttctattcaa tttccggggg ccgtgcggga 240
 30 gtcggggtgg acgtgctctc tctgcccggg ggtcccgaag cgagaggctt cggccctcta 300
 ttggacttca tgtacacttc gcgcctgcgc ctctctccag ccaactgcacc agcagtctta 360
 gcggccgcca cctatttgca gatggagcac gtggtccagg catgccaccg cttcatccag 420
 35 gccagctatg aacctctggg catctccctg cgccccctgg aagcagaacc cccaacaccc 480
 ccaacggccc ctccaccagg tagtcccagg cgctccgaag gacaccaga cccacctact 540
 40 gaatctcgaa gctgcagtca agggcccccc agtccagcca gccctgacct caaggcctgc 600

	aactggaaaa agtacaagta catcgtgcta aactctcagg cctcccaagc agggagcctg	660
	gtcggggaga gaagttcttg tcaaccttgc cccaagcca ggctcccag tggagacgag	720
5	gcctccagca gcagcagcag cagcagcagc agcagcagtg aagaaggacc cattcctggt	780
	ccccagagca ggctctctcc aactgctgcc actgtgcagt tcaaattgtg ggctccagcc	840
10	agtacccct acctcctcac atcccaggct caagacacct ctggatcacc ctctgaacgg	900
	gctcgtccac taccgggagt gaatttttca gctgccagaa ctgtgaggct gtggcagggt	960
	gctcatcggg ggctggactc cttggttctt ggggacgaag acaaacccta taagtgtcag	1020
15	ctgtgccggt cttcgttccg ctacaaggc aaccttgcca gtcaccgtac agtgacaca	1080
	ggggaaaaagc cttaccactg ctcaatctgc ggagcccgtt ttaaccggcc agcaaactg	1140
20	aaaacgcaca gccgcatcca ttcgggagag aagccgtata agtgtgagac gtgcggctcg	1200
	cgctttgtac aggtggcaca tctgcggcg cacgtgctga tccacaccgg ggagaagccc	1260
	tacccttgcc ctacctgcg aaccgcttc cgccacctgc agacctcaa gagccacgtt	1320
25	cgcattcaca ccggagagaa gccttaccac tgcgacctt gtggcctgca tttccggcac	1380
	aagagtcaac tgcggctgca tctgcgccag aaacacggag ctgctaccaa caccaaagtg	1440
30	cactaccaca ttctcgggg gccctagctg agcgcaggcc caggcccccac ttgcttctg	1500
	cgggtgggaa agctgcaggc ccaggccttg cttccctatc aggttgggc ataggggtgt	1560
	gccaggccac tttggtatca gaaattgcca ccctcttaat ttctcactgg ggagagcagg	1620
35	ggtggcagat cctggctaga tctgcctctg ttttgctggt canaccctct tccccacaag	1680
	ccagattgtt tctgaggaga gagctagcta ggggctggga aaggggagag attggagtcc	1740
40	tggctccct aagggaaatag ccctccacct gtggcccca ttgcattcag tttatctgta	1800

aaatataatt tattgaggcc ttggtggc accggggcct tcattcgatt gcatttccca 1860
 ctcccctctt ccacaagtgt gattaaaagt gaccagaaac acagaagggtg agatcacagc 1920
 5 tctgctggca gagattacta gcccttggct ctctcgtttg gcttgggtat tttatattat 1980
 ttctgtcata acttttatct ttagaattgt tctttctcct gtttgtttgc ttgttagttt 2040
 10 gttaaaatg gaaaaagggg ttctctgtgt tctgccctg taattctagg tctggaacct 2100
 ttatttgttc tagggcagct ctggaacat gcgggattgt ggaattgggt caggaacctt 2160
 ctctggtatt ctggatgttg taggttctct agcagtctag aaatggatac agacatttct 2220
 15 ctgttcttca agggtagatg gaaccattat gttgagcca aaatggaagt aataataaat 2280
 gcctcctgga ggctgtgggt gtgggggatt ctgtatctgg attccgtatc actccaactg 2340
 20 gaggctgtgg gtgtggggga ttctgtatct ggattccgta tcaactcaag tggaggctgg 2400
 caggtttttc tgcaagatgg tccagaatct aaaatgtccc attaatctgg tcacttgggt 2460
 ttggctctgc tgtatccatc tatagtggta gagaccacc agggctcaag tggagtccat 2520
 25 catcctcca cgggggcctg ttcttagtac tgagttgatc gctccatggg ggagagatca 2580
 gacattcctt atcagagatg atgtgacctt ttctgactct gccagctctc tatgaatgtt 2640
 30 atggcctagg gaagaatcat gaaactctt agcttgatta gatggtaaacc agtgtaacc 2700
 catcctttac tacagaggca tatgggtttg aatgttacct ggggttctct ctattgagtt 2760
 gagcccttc ttcttttagt gggttttgga catcttctgg caagtgtcca gatgccagaa 2820
 35 ccttcttttc ctctagaagg gatggtgctt ggtaacctta ccttttaaaa gctgggtctg 2880
 tgacctggtc ttcccatccc tgcattcctg tctggaacca gtgaatgcat tagaaccttc 2940
 40 cataggaaaa gaaaaggggc tgagttccat tctgggtttg ctgtagtttg gttgggatta 3000

	ttgttgcat tacagatgta aaagattgac tagcccatag gccaaaggcc tgttctagtt	3060
	gaccaagttt caagtaggat taagagggtt gttgaggggt gcagtttctg gtgtaggcca	3120
5	ggtaggtaga aagtaggaa cagggttgcc tcttggttg gtggagtctc tgaaatgtta	3180
	gaagaagcgc tgaagccttg attgatagtt ctgccccttg ttgccctggg gcttatctga	3240
10	ttatgggacg agggtagaaa gtaagaagca cttttgaatt tgtggggtag aacttcaaca	3300
	ataagtcaagt tctagtggct gtcgcctggg gactagttag aaagctactc ttctccctct	3360
	tccctctttc tcccatggc cccactgcag aattaaagaa ggaagaaggg aaggcggagg	3420
15	agtctataag aaggaatcat gatttctatt tagcagattg gatgggcagg tggagaatgc	3480
	ctgggggtag aaatgttaga tcttgcaaca tcagatcctt ggaataaaga agcctctctg	3540
20	cgcaaaaaaa aaaaaaaaaa aaaa	3564

	<210> 2
	<211> 1440
25	<212> DNA
	<213> Homo sapiens

	<400> 2	
	atgggttccc ccgccgcccc ggaggagcg ctgggctacg tccgcgagtt cactcgccac	60
30	tcctccgacg tgctgggcaa cctcaacgag ctgcgcctgc gcgggacct cactgacgtc	120
	acgctgctgg ttggcgggca acccctcaga gcacacaagg cagttctcat cgctgcagt	180
35	ggcttcttct attcaatttt ccggggccgt gcgggagtcg gggaggacgt gctctctctg	240
	cccgggggtc ccgaagcgag aggccttcgcc cctctattgg acttcatgta cacttcgcgc	300
40	ctgcgcctct ctccagccac tgcaccagca gtcttagcgg ccgccaccta tttgcagatg	360

	gagcacgtgg tccaggcatg ccaccgcttc atccaggcca gctatgaacc tctgggcatc	420
	tccctgcgcc ccctggaagc agaaccccca acaccccaa cggcccctcc accaggtagt	480
5	cccaggcgct ccgaaggaca cccagacca cctactgaat ctcgaagctg cagtcaaggc	540
	ccccccagtc cagccagccc tgacccaag gcctgcaact ggaaaaagta caagtacatc	600
	gtgctaaact ctcaggcctc ccaagcaggg agcctggctg gggagagaag ttctggtaa	660
10	ccttgcccc aagccaggct cccagtgga gacgaggcct ccagcagcag cagcagcagc	720
	agcagcagca gtgaagaagg acccattcct ggtccccaga gcaggctctc tccaactgct	780
15	gccactgtgc agttcaaag tgaggctcca gccagtacc cctacctct ccatcccag	840
	gctcaagaca cctctggatc accctctgaa cgggctcgtc cactaccggg aagtgaattt	900
	ttcagctgcc agaactgtga ggctgtggca ggggtctcat cggggctgga ctccttggtt	960
20	cctggggacg aagacaaacc ctataagtgt cagctgtgcc ggtcttcgtt ccgctacaag	1020
	ggcaaccttg ccagtcacg tacagtgcac acaggggaaa agccttacca ctgctcaatc	1080
25	tgcgagagccc gttttaaccg gccagcaaac ctgaaaacgc acagccgcat ccattcgga	1140
	gagaagccgt ataagtgtga gacgtgcggc tcgcgcttg tacagggtggc acatctgcgg	1200
	gcgcacgtgc tgatccacac cggggagaag ccctaccctt gcctacctg cggaaccgc	1260
30	ttccgccacc tgcagacct caagagccac gttcgcaccc acaccggaga gaagccttac	1320
	cactgcgacc cctgtggcct gcatttcgg cacaagagtc aactgcggct gcatctgcgc	1380
35	cagaaacacg gagctgctac caacacaaa gtgcactacc acattctcg ggggccctag	1440

<210> 3

<211> 65

40 <212> DNA

<213> Homo sapiens

 <400> 3
 ataccgga ctcctaagc cttctattag ctccaataat agtaagcctg tcgaagacaa 60
 5 agatg 65

 <210> 4
 10 <211> 70
 <212> DNA
 <213> Homo sapiens

 <400> 4
 15 gcctgtgtcc cctagactcc aactcagcaa cggaataga actctgaccc tgtttaacgt 60
 gaccaggaac 70

 20 <210> 5
 <211> 70
 <212> DNA
 <213> Homo sapiens

 25 <400> 5
 acgtgcttta cggacccgat gctcctacaa tcagccctct aaacacaagc tatagatcag 60
 gggaaaatct 70

 30 <210> 6
 <211> 70
 <212> DNA
 <213> Homo sapiens
 35 <400> 6
 acgttaaaca gggtcagagt tctatttccg ttgctgagtt ggagtctagg ggacacaggc 60
 agggactggt 70
 40

<210> 7
 <211> 70
 <212> DNA
 5 <213> Homo sapiens

 <400> 7
 ctgatctata gcttgtgttt agagggctga ttgtaggagc atcgggtccg taaagcacgt 60
 10 tgagaatcac 70

 <210> 8
 <211> 63
 15 <212> DNA
 <213> Homo sapiens

 <400> 8
 gatccactat tgttcacggt aatattggga atgaacagtt cctgggtgga ctggttgaaa 60
 20 gtg 63

 <210> 9
 25 <211> 70
 <212> DNA
 <213> Homo sapiens

 <400> 9
 30 gacacagcaa gctacaaatg cgaaacccaa aatccagtca gcgccaggag gtctgattca 60
 gtgattctca 70

 35 <210> 10
 <211> 70
 <212> DNA
 <213> Homo sapiens

 40 <400> 10

	tgaatcagac ctcctggcgc tgactggatt ttgggtttcg catttgtagc ttgctgtgtc	60
	gttcctggtc	70
5	<210> 11 <211> 79 <212> DNA <213> Homo sapiens	
10	<400> 11 gatcctacac gtgccaagct cacaatagcg acaccggact caaccgcaca accgtgacga	60
	cgattaccgt gtatgccga	79
15	<210> 12 <211> 70 <212> DNA <213> Homo sapiens	
20	<400> 12 catcctcaac tgggttagaa ttgttactag ttatgaatgg ttttggtggc tcggcataca	60
25	cggtaatcgt	70
	<210> 13 <211> 80 <212> DNA <213> Homo sapiens	
30	<400> 13 ttctaacca gttgaggatg aggacgcagt tgcattaact tgtgagccag agattcaaaa	60
35	taccacttat ttatgggtggg	80
	<210> 14 <211> 80	
40		

<212> DNA

<213> Homo sapiens

<400> 14

5 gtctaataatgat aaccgcacat tgacactcct gtccggttact cgcaatgatg taggacctta 60

tgagtgtggc attcagaatg 80

10 <210> 15

<211> 80

<212> DNA

<213> Homo sapiens

15 <400> 15

tttgtatggc ccagacgacc caactatatc tccatcatac acctactacc gtcccggcgt 60

gaacttgagc ctttcttgcc 80

20

<210> 16

<211> 80

<212> DNA

<213> Homo sapiens

25

<400> 16

tgatggaaac attcagcagc atactcaaga gttatttata agcaacataa ctgagaagaa 60

cagcggactc tatacttgcc 80

30

<210> 17

<211> 80

<212> DNA

35 <213> Homo sapiens

<400> 17

taaaacaata actgtttccg cgagagctgcc caagccctcc atctccagca acaactccaa 60

40 acccgtggag gacaaggatg 80

<210> 18
 <211> 80
 5 <212> DNA
 <213> Homo sapiens

 <400> 18
 atgtgcggtt atcattagac aactgcaagc gtgggctaac cggcaaactt tggttattga 60
 10
 cccaccataa ataagtggta 80

 <210> 19
 15 <211> 80
 <212> DNA
 <213> Homo sapiens

 <400> 19
 20 ggtcgtctgg gccatacaaa acattaagga taacagggtc ggagtgatca acggataatt 60

 cattctgaat gccacactca 80

 25 <210> 20
 <211> 80
 <212> DNA
 <213> Homo sapiens

 30 <400> 20
 gctgctgaat gtttccatca atcagccagg agtactgtgc agggggggtg gatgctgcat 60

 ggcaagaaag gctcaagttc 80

 35
 <210> 21
 <211> 80
 <212> DNA
 <213> Homo sapiens
 40

<400> 21
 cggaacagt tattgtttta actgtagtcc tgctgtgacc actggctgag ttattggcct 60

 ggcaagtata gagtccgctg 80
 5

 <210> 22
 <211> 47
 <212> DNA
 10 <213> Homo sapiens

 <400> 22
 cctcaggttc acaggtgaag gccacagcat cctgttcctc cacgggt 47

 15
 <210> 23
 <211> 2106
 <212> DNA
 <213> Homo sapiens
 20
 <400> 23
 atggagtctc cctcggcccc tccccacaga tgggtgcatcc cctggcagag gtcctgctc 60

 acagcctcac ttctaacctt ctggaacccg cccaccactg ccaagctcac tattgaatcc 120
 25
 acgccgttca atgtcgcaga ggggaaggag gtgcttctac ttgtccacaa tctgccccag 180

 catctttttg gctacagctg gtacaaagggt gaaagagtgg atggcaaccg tcaaattata 240
 30
 ggatatgtaa taggaactca acaagctacc ccagggcccg catacagtgg tcgagagata 300

 atatacccca atgcatccct gctgatccag aacatcatcc agaatgacac aggattctac 360

 accctacag tcataaagtc agatcttggt aatgaagaag caactggcca gttccgggta 420
 35
 taccgggagc tgcccaagcc ctccatctcc agcaacaact ccaaaccgt ggaggacaag 480

 gatgctgtgg ccttcacctg tgaacctgag actcaggacg caacctacct gtggtgggta 540
 40
 aacaatcaga gcctcccggt cagtcccagg ctgcagctgt ccaatggcaa caggaccctc 600

actctattca atgtcacaag aaatgacaca gcaagctaca aatgtgaaac ccagaaccca 660
 gtgagtgcc a ggcgagtgga ttcatgtatc ctgaatgtcc tctatggccc ggatgcccc 720
 5 accatttccc ctctaaacac atcttacaga tcaggggaaa atctgaacct ctctgcccac 780
 gcagcctcta acccacctgc acagtactct tggtttgtca atgggacttt ccagcaatcc 840
 10 acccaagagc tctttatccc caacatcact gtgaataata gtggatccta tacgtgcca 900
 gcccataact cagacactgg cctcaatagg accacagtca cgacgatcac agtctatgag 960
 ccacccaac cttcatcac cagcaacaac tccaaccccg tggaggatga ggatgctgta 1020
 15 gccttaacct gtgaacctga gattcagaac acaacctacc tgtgggtggg aaataatcag 1080
 agcctcccg tcagtcaccg gctgcagctg tccaatgaca acaggaccct cactctactc 1140
 20 agtgtcaca ggaatgatgt aggaccctat gagtgtggaa tccagaacga attaagtgtt 1200
 gaccacagcg acccagtcac cctgaatgtc ctctatggcc cagacgacc caccatttcc 1260
 ccctcataca cctattaccg tccaggggtg aacctcagcc tctcctgcca tgcagcctct 1320
 25 aacccacctg cacagtattc ttggctgatt gatgggaaca tccagcaaca cacacaagag 1380
 ctctttatct ccaacatcac tgagaagaac agcggactct atacctgcca ggccaataac 1440
 30 tcagccagtg gccacagcag gactacagtc aagacaatca cagtctctgc ggagctgccc 1500
 aagccctcca tctccagca caactccaaa cccgtggagg acaaggatgc tgtggccttc 1560
 acctgtgaac ctgaggtca gaacacaacc tacctgtggt gggtaatgg tcagagcctc 1620
 35 ccagtcagtc ccaggctgca gctgtccaat ggcaacagga cctcactct attcaatgtc 1680
 acaagaaatg acgcaagagc ctatgtatgt ggaatccaga actcagtgag tgcaaaccgc 1740
 40 agtgaccag tcaccctgga tgcctctat gggccggaca ccccatcat ttcccccca 1800

gactcgtctt acctttcggg agcggacctc aacctctcct gccactcggc ctctaaccca 1860

tccccgcagt attcttggcg tatcaatggg ataccgcagc aacacacaca agttctcttt 1920

5 atcgccaaaa tcacgccaaa taataacggg acctatgcct gttttgtctc taacttggct 1980

actggccgca ataattccat agtcaagagc atcacagtct ctgcatctgg aacttctcct 2040

10 ggtctctcag ctggggccac tgcgggcac atgattggag tgctggttgg ggttgctctg 2100

atatag 2106

15 <210> 24
 <211> 47
 <212> DNA
 <213> synthetic

20 <400> 24
 ggacggtagt aggtgtatga tggagatata gttgggtcgt ctgggcc 47

<210> 25
 25 <211> 27
 <212> DNA
 <213> Synthetic

<400> 25
 30 cagaatgaat tatccgttga tcactcc 27

<210> 26
 <211> 45
 35 <212> DNA
 <213> Synthetic

<400> 26
 cgtgacgacg attaccgtgt atgagccacc aaaaccattc ataac 45

40

<210> 27
 <211> 45
 <212> DNA
 5 <213> Synthetic

<400> 27
 gttatgaatg gttttggtgg ctcatcacg gtaatcgtcg tcacg 45

10
 <210> 28
 <211> 2106
 <212> DNA
 <213> Synthetic

15
 <400> 28
 atggagtctc cctcggtccc tccccacaga tggatcatcc cctggcagag gctcctgctc 60

20
 acagcctcac ttctaacctt ctggaacccg cccaccactg ccaagctcac tattgaatcc 120
 acgccgttca atgtcgcaga ggggaaggag gtgcttctac ttgtccacaa tctgccccag 180
 catctttttg gctacagctg gtacaaaggt gaaagagtgg atggcaaccg tcaaattata 240

25
 ggatatgtaa taggaactca acaagctacc ccagggtccc catacagtgg tcgagagata 300
 atatacccca atgcatccct gctgatccag aacatcatcc agaatgacac aggattctac 360

30
 accctacacg tcataaagtc agatcttgtg aatgaagaag caactggcca gttccgggta 420
 taccgggaac tccctaagcc ttctattagc tccaataata gtaagcctgt cgaagacaaa 480
 gatgccgtcg cttttacatg cgagcccgaa actcaagacg caacatatct ctggtgggtg 540

35
 aacaaccagt ccctgcctgt gtcccctaga ctccaactca gcaacggaaa tagaactctg 600
 accctgttta acgtgaccag gaacgacaca gcaagctaca aatgcgaaac ccaaaatcca 660
 gtcagcgcca ggaggtctga ttcaagtatt ctcaacgtgc ttacggacc cgatgctcct 720

40

acaatcagcc ctctaaacac aagctataga tcaggggaaa atctgaatct gagctgtcat 780
 gccgctagca atcctcccg ccaatacagc tggtttgta atggcacttt ccaacagtcc 840
 5 acccaggaac tgttcattcc caatattacc gtgaacaata gtggatccta cacgtgccaa 900
 gctcacaata gcgacaccgg actcaaccgc acaaccgtga cgacgattac cgtgtatgag 960
 ccacaaaac cattcataac tagtaacaat tctaaccag ttgaggatga ggacgcagtt 1020
 10 gcattaactt gtgagccaga gattcaaaat accacttatt tatggtgggt caataaccaa 1080
 agtttgccgg ttagcccagc cttgcagttg tctaataata accgcacatt gacactcctg 1140
 15 tccgttactc gcaatgatgt aggaccttat gagtgtggca ttcagaatga attatccgtt 1200
 gatcactccg accctgttat ccttaatgtt ttgtatggcc cagacgaccc aactatatct 1260
 ccatcataca cctactaccg tcccggcgtg aacttgagcc tttcttgcca tgcagcatcc 1320
 20 aacccccctg cacagtactc ctggctgatt gatggaaaca ttcagcagca tactcaagag 1380
 ttatttataa gcaacataac tgagaagaac agcggactct atacttgcca ggccaataac 1440
 25 tcagccagtg gtcacagcag gactacagtt aaaacaataa ctgtttccgc ggagctgccc 1500
 aagccctcca tctccagcaa caactccaaa cccgtggagg acaaggatgc tgtggccttc 1560
 acctgtgaac ctgaggctca gaacacaacc tacctgtggt gggtaaatgg tcagagcctc 1620
 30 ccagtcagtc ccaggctgca gctgtccaat ggcaacagga ccctcactct attcaatgtc 1680
 acaagaaatg acgcaagagc ctatgtatgt ggaatccaga actcagtgag tgcaaaccgc 1740
 35 agtgacccag tcaccctgga tgtcctctat gggccggaca ccccatcat ttcccccca 1800
 gactcgtctt accttcggg agcggacctc aacctctcct gccactcggc ctctaacca 1860
 tccccgcagt attcttggcg tatcaatggg ataccgcagc aacacacaca agttctcttt 1920
 40

atcgccaaaa tcacgccaaa taataacggg acctatgcct gttttgtctc taacttggct 1980
 actggccgca ataattccat agtcaagagc atcacagtct ctgcatctgg aacttctcct 2040
 5 ggtctctcag ctggggccac tgcggcatc atgattggag tgctggttg ggttgctctg 2100
 atatag 2106

 10 <210> 29
 <211> 35
 <212> DNA
 <213> Synthetic

 15 <400> 29
 ctggcgcgcc ttctttattc tataacttaaa aagtg 35

 <210> 30
 20 <211> 36
 <212> DNA
 <213> Synthetic

 <400> 30
 25 ctggtaccag aaaaactata tcagagcaac cccaac 36

 <210> 31
 <211> 26
 30 <212> DNA
 <213> Synthetic

 <400> 31
 ttggatccat ggagtctccc tcggcc 26
 35

 <210> 32
 <211> 26
 <212> DNA
 40 <213> Synthetic

<400> 32

ttggatccct atatcagagc aacccc

26